

WATERSHED CHARACTERIZATION

| | | |
|-----------|--|----------|
| B1 | METHODOLOGY AND GAPS | 1 |
| B1.1 | Data Resources Matrix | 1 |
| B1.2 | Monitoring Data Sources | 1 |
| B1.2.1 | Stream Flow Gauging | 5 |
| B1.2.2 | Precipitation/Meteorological Gauging | 6 |
| B1.2.3 | Snow Cover Monitoring | 7 |
| B1.2.4 | Groundwater Monitoring..... | 7 |
| B1.2.5 | Surface Water Quality | 8 |
| B1.2.6 | Low-Flow Stream Flow Surveys..... | 10 |
| B1.2.7 | Biological Monitoring | 12 |
| B1.2.8 | Coastal Wetland Monitoring..... | 12 |
| B1.3 | Information Management System..... | 16 |
| B1.4 | Methods of Analysis | 16 |
| B1.5 | Surface Water Quality Data Analysis and Reporting..... | 17 |
| B1.5.1 | Exploratory Analysis | 17 |
| B1.5.2 | Statistical Analysis | 17 |
| B1.5.3 | Reporting Results | 17 |
| B1.6 | Groundwater Quality Data Analysis and Reporting | 26 |
| B1.6.1 | Data Compilation | 26 |
| B1.6.2 | Data Analysis | 26 |
| B1.6.3 | Analysis of Trends at Each Monitoring Well | 26 |
| B1.6.4 | Aquifer Characterization | 33 |
| B1.7 | Limitations: Data, Assumptions, and Methods | 33 |
| B1.7.1 | Data | 33 |
| B1.7.2 | Addressing Data | 34 |
| B1.7.3 | Assumptions..... | 34 |
| B1.7.4 | Method Limitations..... | 35 |

FIGURES

| | |
|---|----|
| Figure B-1: CLOSPA Water Monitoring Network — 2006..... | 4 |
| Figure B-2: Ontario Benthic Biomonitoring Network (OBBN) within the Study Area..... | 13 |
| Figure B-3: Coastal Wetland Monitoring Project Locations, Durham Region..... | 14 |

TABLES

| | |
|---|----|
| Table B-1: Monitoring Databases and Data Descriptions | 3 |
| Table B-2: Surface Water Gauge Details, CLOSPA/Provincial Stream Gauge Network Stations | 5 |
| Table B-3: Meteorological Station Details, CLOSPA and AES Stations..... | 6 |
| Table B-4: Snow Course Station Details, CLOSPA Stations | 7 |
| Table B-5: PGMN Groundwater Monitoring Well Details..... | 8 |
| Table B-6: Surface Water Quality Station Details, PWQMN and CLOSPA Stations | 10 |

Table B-7: Low-Flow Monitoring Details, CLOSPA Stations 11
Table B-8: Durham Region Coastal Wetlands Currently Monitored..... 15
Table B-9: Statistical Results for Select Surface Water Quality Sites within CLOSPA 25
Table B-10: List of all Groundwater Quality Samples within CLOSPA with Parameters that have
Exceeded the ODWS at Least once..... 32
Table B-11: Data Identified 34

B1 METHODOLOGY AND GAPS

B1.1 DATA RESOURCES MATRIX

To organize the data sources required for the drinking water source protection (SWP) watershed characterization, the Province has developed an Excel file called the SWP Data Requirements Matrix. The matrix is intended to:

- Provide a complete list of available data sets for SWP;
- Help inventory and evaluate local data;
- Help identify data gaps;
- Facilitate data request process; and
- Facilitate communications around data between neighbouring Conservation Authorities and their SWP watershed region.

The matrix includes data set names, data descriptions, data access, data sources, and links to metadata. The file also includes a list of data sources required to build particular maps. Requests for data have been made by Central Lake Ontario Source Water Protection Area (CLOSPA) through SWP activities to both the Province and local municipalities. An inventory of the data and metadata received to date is being maintained by Central Lake Ontario Conservation Authority (CLOCA).

B1.2 MONITORING DATA SOURCES

CLOSPA's monitoring networks provide an ongoing source of data sets that support numerous programs, including SWP planning. The source protection process involves the following steps:

- Developing an understanding of the flow system (surface water and groundwater);
- Cataloguing the various potential contaminant sources; and
- Assessing the risk that these potential contaminants pose to the water supply resources on a watershed basis.

CLOSPA's monitoring databases that are relevant to source water protection planning are listed in **Table B-1** which includes data type, status, and spatial coverage. CLOSPA's monitoring network incorporates both provincial and federal monitoring partnership programs. This monitoring network collects information pertaining, but not limited to, the following data types:

- Streamflow (stream gauges and low flows);
 - Quantity — Environment Canada HYDAT and Conservation Authority, and
 - Quality — Conservation Authority and Provincial Water Quality Monitoring Network (PWQMN).
- Climate;
- Groundwater; and
 - Quantity — Water levels, and
 - Quality — Provincial Groundwater Monitoring Network (PGMN).
- Groundwater monitoring networks operated by municipal partners related to the municipal well areas.

Other monitoring programs, such as aquatic ecosystem studies conducted by the Province and conservation authorities, also contribute to the development of a SWP plan.

Climate data is collected, mainly by Environment Canada, at the locations shown on **Figure B-1**. These data are augmented by monitoring stations operated by the conservation authorities. Continuous streamflow gauges are also maintained by the federal government (Water Survey of Canada). These locations are also shown on **Table B-2**. Stream water quality is monitored through the Provincial Water Quality Monitoring Network (PWQMN), and is augmented by other stations operated by the conservation authorities to contribute to various programs. These are described further in the watershed characterization report listed in **Chapter 7**.

Streamflow quality is important to the overall monitoring of watershed health, and is necessary to determine chemical loadings to Lake Ontario, the source of water supply for the majority of the population. Groundwater quality and quantity (water levels) are monitored by the Province and the conservation authorities at the locations shown on **Figure B-1**. Groundwater levels and quality monitoring is also conducted by municipal partners with locations mainly associated with the wellfield areas.

The information obtained from the monitoring network is applied in the various analyses that have or are being conducted as prescribed in the *Technical Rules*.

| Database Name | Data | | | | |
|--|--|--------|-------------------------------------|--------------------------|--|
| | Type | Format | Period of Record | Coverage Area | Recording/Collection Frequency |
| Durham Region Coastal Wetlands Monitoring Database | Turbidity | access | 2002 – present | 8 coastal wetlands | Monthly readings |
| | Water levels | excel | 2003 – present | 5 coastal wetlands | Continuous readings |
| | Water temperature | excel | 2003 – present | 5 coastal wetlands | Continuous readings |
| | Sediment quality | access | 2002 | 8 coastal wetlands | 5-year rotation collection |
| | Fish community | access | 2003 – present | 8 coastal wetlands | Monthly collection |
| | Invertebrate | access | 2003 – present | 8 coastal wetlands | Monthly collection |
| | Submerged plants | access | 2003 – present | 8 coastal wetlands | Monthly collection |
| Wetland Evaluation Database | MNR evaluation reports | paper | 2005 | 15 wetlands | 5 year rotation |
| ARMP Bio-Monitoring Database | Water Quality Index values (WQI), status and system type | excel | 1996 – 2004 (terminated) | ARMPs per watershed | One collection per site per ARMP |
| | Water temperature | excel | 1996 – 2004 (terminated) | ARMPs per watershed | One collection per site per ARMP |
| OBBN Bio-Monitoring Database | TBD - Reference Condition Approach (RCA) | TBD | Initiated 2005 | OBBN sites TBD | TBD |
| | TBD - Stream morphology | TBD | Initiated 2005 | OBBN sites TBD | TBD |
| Species Database | Terrestrial species attributes | access | 2003 – present | Jurisdiction | Seasonal collection |
| Water Monitoring Network Databases | Groundwater quality (CLOSPA/PGMN) | access | 2001 – present | 16 sites | 2 samples collected per site per year |
| | Groundwater static measurements (CLOSPA/PGMN) | access | 2003 – present | 16 sites | 2 readings per site per year |
| | Groundwater continuous levels (CLOSPA/PGMN) | access | 2001 – present | 16 sites | Continuous readings |
| | Surface water quality (CLOSPA/PWQMN) | access | 1965 – present | 19 sites | Monthly collection at PWQMN sites; two samples collected per year at CLOSPA sites; |
| | Surface water levels | access | 1959 – present depending on station | 10 locations + 3 in 2005 | Continuous |
| | Surface water flows | access | 1959 – present depending on station | 5 locations | Continuous |

| Database Name | Data | | | | |
|-----------------------|--|----------------|------------------|-----------------|--------------------------------------|
| | Type | Format | Period of Record | Coverage Area | Recording/Collection Frequency |
| | Rainfall | access | 1999 – present | 7 locations | Continuous |
| | Snow pack | access | 1980 – present | 4 locations | Bi-monthly - seasonal |
| | Humidity | access | 2001 – present | 2 locations | Continuous |
| | Barometric pressure | access | 2001 – present | 2 locations | Continuous |
| | Wind speed/direction | access | 2001 – present | 2 locations | Continuous |
| | Air temperature | access | 2001 – present | 7 locations | Continuous |
| | Rainfall | access | 1959 – present | 7 locations | Continuous |
| | Low flows (CLOSPA) | access | 2000 - present | 67 sites | 4 – 6 measurements per site per year |
| | Low flows (ORMGP) | e:DAT | 2002 | 46 sites | 1 measurement per site |
| | Stream morphology | e:DAT | 2002 | 46 sites | 1 measurement per site |
| | Site locations | access | Current | All sites | As added/removed |
| Field notes | excel | 2001 - present | Most sites | As required | |
| PTTW Database | Potential contaminant threats, locations, and attributes | TBD | 2002 - present | 83 active sites | As Identified |
| ORMGP Database | Subsurface/well data | access | 1950s – present | Jurisdiction | As identified |
| | Climatic data | access | 1960s – present | Jurisdiction | As identified |
| | Surface water data | access | 1960s – present | Jurisdiction | As identified |
| | PTTW data | access | 2002 – present | Jurisdiction | As identified |

Table B-1: Monitoring Databases and Data Descriptions




Figure B-1: CLOSPA Water Monitoring Network — 2006

B1.2.1 Stream Flow Gauging

Stream gauge data are required for water budgets, assimilative capacity studies, water takings, aquatic studies, and recharge and discharge analyses. Total flows, base flows, mean daily flows, and mean monthly flow information is derived from the raw level data and stream section and profile survey information.

There are 13 active stations (Table B-2), which include both Environment Canada’s Water Survey of Canada (WSC) and CLOSPA stations. Active gauge stations do not exist on Black, Farewell, Corbett, Goodman, Robinson, Tooley, Darlington, Westside, and Bennett creeks. Historical records exist for discontinued WSC gauge stations sited on Farewell and Soper creeks. Additional stations will be required to generate the data required for SWP planning initiatives.

| STATION ID | WATERSHED | LOCATION | PARAMETER | RECORD FREQUENCY | PERIOD OF RECORD |
|-------------------------|---------------------------------|---------------|-------------|------------------|---------------------------|
| 02HC018 (WSC) | Lynde Creek (main branch) | Dundas St. | Water Level | hr | 1959–current |
| | | | Rainfall | | |
| 02HD008 (WSC) | Oshawa Creek (main branch) | Taunton Rd. | Water Level | hr | 1959–current |
| | | | Rainfall | 15min | |
| 02HD006 (WSC) | Bowmanville Creek (main branch) | Jackman Rd. | Water level | hr | 1959–current |
| | | | Rainfall | | Proposed–2005 |
| 02HC055 (WSC) | Lynde Creek (west branch) | Kinsale | Water Level | hr | 2002–current |
| | | | Rainfall | | |
| | | | Water Temp | | |
| | | | Air Temp | | |
| 02HC054 (WSC) | Lynde Creek (east Branch) | Brooklin | Water Level | hr | 2002–current |
| | | | Water Temp | | |
| Hampton | Bowmanville Creek (west branch) | Hampton | Water Level | hr | 2003–current |
| | | | Rainfall | | |
| | | | Water Temp | | |
| | | | Air Temp | | |
| Bowtaunt | Bowmanville Creek (east branch) | Taunton Rd. | Water Level | hr | Installation not complete |
| | | | Water Temp | | |
| OshWest | Oshawa Creek (west branch) | Conlin Rd. | Water Level | hr | 2001–current |
| | | | Air Temp | | |
| OshEast | Oshawa Creek (east branch) | Conlin Rd. | Water Level | hr | 2001–current |
| | | | Air Temp | | |
| OshMain | Oshawa Creek (main branch) | Thomas St. | Water Level | hr | 2001–current |
| | | | Air Temp | | |
| Pringle | Pringle Creek (main branch) | Consumers Rd. | Water Level | hr | 2000–current |
| SopEast (02HD223) | Soper Creek (east branch) | Taunton Rd. | Water Level | hr | Proposed–2005 |
| SopWest (02HD222) (WSC) | Soper Creek (west Branch) | Taunton Rd. | Water Level | hr | Proposed–2005 |

Table B-2: Surface Water Gauge Details, CLOSPA/Provincial Stream Gauge Network Stations

B1.2.2 Precipitation/Meteorological Gauging

Monitoring and measuring precipitation is a fairly simple process. One must obtain an accurate sample of the precipitation falling at the location of the gauge and have sufficient spatial coverage throughout the watershed to permit accurate estimates of the volume of water falling on a watershed. This information is currently compared with runoff volumes and quantitative hydrologic forecasting. Two types of climate station measurement locations operate within the watershed: five rain gauges operated and maintained by CLOSPA, and ten operated by the Atmospheric Environment Service (AES) (**Table B-3**). The data are collected by AES and are available from the Environment Canada website. Historical precipitation records used for the CLOSPA watershed are primarily recorded at the City of Oshawa WPCP station (6155878), which records mean daily rainfall and snowfall events. Data from AES stations are archived and found at: <https://climate.weather.gc.ca/>

AES data are periodically exported into the Oak Ridges Moraine Groundwater Program database for sites within the study area and are listed in **Table B-3**.

| STATION ID | WATERSHED | LOCATION | PARAMETER | PERIOD OF RECORD |
|---------------|-------------------|----------------------|-----------------|------------------|
| Prec1 | Oshawa Creek | Coates Rd. | Rainfall | 1999–present |
| | | | Temperature | 2002-present |
| | | | Wind | |
| | | | Speed/Direction | |
| | | | BP | |
| | | | Humidity | |
| Prec2 | Oshawa Creek | Howden Rd. | Rainfall | 1999-present |
| Prec3 | Oshawa Creek | Whiting Ave. | Rainfall | 1999-present |
| | | | Temperature | 2001-present |
| | | | Wind | |
| | | | Speed/Direction | |
| | | | BP | |
| | | | Humidity | |
| Prec4 | Lynde Creek | Chalk Lake | Climate Stn. | 2003-present |
| Prec5 | Soper Creek | Woodley Rd. | | 2003-present |
| 02HC018 (WSC) | Lynde Creek | Dundas St. | | Current |
| 20HD008 (WSC) | Oshawa Creek | Taunton Rd. | | Current |
| 6150830 (AES) | Soper Creek | Mostert | | Historical |
| 6151042 (AES) | Bowmanville Creek | Burketon McLaughlin | | Historical |
| 6154611 (AES) | Bowmanville Creek | Long Sault IHD | | Historical |
| 6155874 (AES) | Pringle Creek | Whitby WPCP | | Historical |
| 6155876 (AES) | Oshawa Creek | City of Oshawa | | Historical |
| 6155877 (AES) | Harmony Creek | Oshawa Fire Hall | | Current |
| 6155878 (AES) | Oshawa Creek | Oshawa WPCP | Current | |
| 6156634 (AES) | Bennett Creek | Port Darlington WCPC | Current | |
| 6157884 (AES) | Soper Creek | Soper Creek WPCP | Current | |
| 6159048 (AES) | Bowmanville Creek | Tyrone | Historical | |

Table B-3: Meteorological Station Details, CLOSPA and AES Stations

B1.2.3 Snow Cover Monitoring

CLOSPA operates and maintains five snow course survey locations in the watershed (Table B-4), located in the Oak Ridges Moraine and Iroquois Beach physiographic regions. A snow course consists of a series of numbered posts driven into the ground 30 metres apart, usually in a straight line. The water is calculated, based on the weight of the snow in a core sampler. One ounce of snow in the sampler contains the equivalent of one inch of water. Snow course measurements are taken twice monthly, from December to May.

| STATION ID | WATERSHED | LOCATION | PARAMETER | RECORD FREQUENCY (Nov-May) | PERIOD OF RECORD |
|------------|-------------------|--------------------|-----------|----------------------------|------------------|
| 4701 | Lynde Creek | Heron Rd. | Snow | 2/month | current |
| 4909 | Lynde Creek | Coronation Rd. | | | |
| 4802 | Oshawa Creek | Coates Rd. | | | |
| 4903 | Bowmanville Creek | Woodley Rd. | | | |
| 4902 | Soper Creek | Stephen's Hill Rd. | | | |

Table B-4: Snow Course Station Details, CLOSPA Stations

B1.2.4 Groundwater Monitoring

In partnership with the Ministry of the Environment and Climate Change (MOECC), CLOSPA operates and maintains a network of 16 groundwater monitoring wells located throughout the watershed (Table B-5). Loggers were installed in the monitoring wells from 2000 to 2003 and automatically record water levels and temperature. Dataloggers measure absolute pressure (water pressure + atmospheric pressure), expressed in centimetres of water column.

The data are downloaded and sent electronically to the MOECC Provincial Groundwater Monitoring Information System database (PGMIS). The data are locally exported from PGMIS into the YDPT database using a SITEFX (specialized software) interface. CLOSPA is required to perform QA/QC activities to verify the continued accuracy of the data. Water levels are periodically measured manually to ensure that the automated systems are functioning correctly. QA/QC activities for all CLOSPA wells have not been completed at this time. Efforts are being made to align non-SWP funded program deliverables to support SWP analytical requirements.

| STATION ID | WATERSHED | LOCATION | PERIOD OF RECORD | PARAMETER | RECORD FREQUENCY |
|-------------|-------------------|-------------------|------------------|-------------|------------------|
| W0000040-1 | Farewell Creek | Near Courtice Rd. | 2001 – present | Water level | hr |
| W0000042-1 | Soper Creek | Region Rd. 20 | | | |
| W0000043-3 | Bowmanville Creek | Fourth St. | | | |
| W0000044-2 | Soper Creek | Bethesda Rd. | | | |
| W0000044-3 | | | | | |
| W0000049-1 | Oshawa Creek | Raglan Rd. | | | |
| W00000167-1 | Bowmanville Creek | Holt Rd. | 2002 – present | | |
| W00000166-1 | | | | | |
| W00000263-1 | Lynde Creek | Middle March Rd. | | | |
| W00000262-1 | Oshawa Creek | Grass Grove Rd. | | | |
| W00000261-1 | Lynde Creek | Coronation Rd. | | | |
| W00000264-1 | Bowmanville Creek | Grasshopper Rd. | | | |
| W00000264-2 | | | | | |
| W00000265-1 | | | | | |

Table B-5: PGMN Groundwater Monitoring Well Details

In general, sites monitor significant groundwater recharge areas within the Oak Ridges Moraine and in watersheds that originate from the Lake Iroquois Beach. All wells but one monitor surficial formations.

Water samples are also collected from each well twice a year and are analyzed routinely for general chemistry and metals. Every fourth or fifth year, samples are collected at certain locations and analyzed for volatile organics and/or pesticides. This sampling frequency may be adjusted depending on monitor location, interval, local land use, or other identified contaminant issue. Historical water level and quality data are available in hard copy from wells monitored by the MOECC through the IHD studies undertaken through the 1960s and intermittently into the 1980s.

Groundwater level data collected from listed monitoring wells is locally exported from the provincial PGMIS database into the local ORMGP database for use in analysis. Results of electronic water quality analysis from the laboratory are currently input directly into the local ORMGP database.

B1.2.5 Surface Water Quality

Chemical and physical characteristics of surface water quality in CLOSPA are monitored through the Provincial Surface Water Quality Monitoring Network (PWQMN). CLOSPA participates in this program by collecting monthly samples from April through November. The samples are analyzed for a range of water quality indicators, including temperature, pH, conductivity, turbidity, suspended solids, major ions, nutrients, metals, and pesticides, in order to screen overall water quality. CLOSPA currently monitors nine PWQMN stations located at watershed and subwatershed outlets (**Table B-6**).

Historical data sets have existed for each site extending back to the early 1960s, though significant gaps in the dataset have been identified. In addition to the sites currently monitored, there are eight historical PWQMN sites that also support long-term trend analysis.

| Current Sites | | | | | | | | |
|----------------|-----------|---------------------------|------------------------------|-------|------|------------|-----------|----------------|
| MOE Station ID | CLOSPA ID | Creek/Watershed | Location | First | Last | Re-started | Frequency | Program |
| 6010800102 | 1 | Lynde Creek | Victoria St, Whitby | 1964 | 1997 | 2003 | 8/year | PWQMN / CLOSPA |
| 6011100102 | 2 | Oshawa Creek | Simcoe St. South, Oshawa | 1964 | 1997 | 2003 | | |
| 6011200302 | 3 | Farewell Creek | Colonel Sam Drive, Oshawa | 1980 | 1997 | 2003 | | |
| 6011600102 | 4 | Bowmanville Creek | West Beach Rd, Bowmanville | 1964 | 1997 | 2003 | | |
| 6010800402 | 8 | Lynde Creek | Baldwin St, Brooklin | 1977 | 1994 | 2003 | | |
| 6011100302 | 10 | Oshawa Creek | Conlin Road, Oshawa | 2003 | | | | |
| 6011200502 | 14 | Black Creek | Trulls Road, Courtice | 2003 | | | | |
| 6011600502 | 15 | Bowmanville Creek | Hampton Conservation Area | 2003 | | | | |
| 6011600602 | 17 | Bowmanville Creek | Long Sault Conservation Area | 2003 | | | | |
| 6011600202 | 5 | Soper Creek | West Beach Rd, Bowmanville | 1967 | 1994 | 2003 | 2/year | CLOSPA |
| | 9 | Lynde Creek | Heber Down Conservation Area | 2004 | | | | |
| | 11 | Oshawa Creek | Conlin Road, Oshawa | 2004 | | | | |
| 6011200102 | 12 | Harmony Creek | Bloor St, Oshawa | 1964 | 1981 | 2005 | | |
| | 13 | Farewell Creek | Nash Road, Courtice | 2005 | | | | |
| | 16 | Bowmanville Creek | Taunton Rd, Clarington | 2004 | | | | |
| | 18 | Soper Creek (west branch) | Taunton Rd, Clarington | 2004 | | | | |
| | 19 | Soper Creek (east branch) | Taunton Rd, Clarington | 2004 | | | | |
| | 20 | Soper Creek (east branch) | Gibbs Rd north Conc. 7 | 2004 | | | | |
| | 21 | Soper Creek (east branch) | Lambs Road, Clarington | 2005 | | | | |

| Historical Sites | | | | | | | | |
|------------------|-----------|------------------|-------------------------------|-------|------|------------|-----------|---------|
| MOE Station ID | CLOSPA ID | Creek/Watershed | Location | First | Last | Re-started | Frequency | Program |
| 06010800202 | 30 | Lynde Creek | Baldwin St, N of Taunton Rd | 1977 | 1978 | | | |
| 06010800302 | 31 | Lynde Creek | Winchester Rd, E of Hwy 7/12 | 1977 | 1988 | | | |
| 06010900102 | 32 | Pringle Creek | Brock St, Whitby | 1964 | 1987 | | | |
| 06010900302 | 33 | Pringle Creek | Watson St, Whitby | 1972 | 1994 | | | |
| 06011100202 | 34 | Montgomery Creek | Harbour Rd, Oshawa | 1966 | 1994 | | | |
| 06011600302 | 35 | Soper Creek | King St E, Hwy 2, Bowmanville | 1968 | 1990 | | | |

Table B-6: Surface Water Quality Station Details, PWQMN and CLOSPA Stations

B1.2.6 Low-Flow Stream Flow Surveys

CLOSPA is working with the Ministry of Natural Resources and Forestry, and the MOECC on the Low Water Response Program. This program monitors rainfall and streamflow within the creeks of CLOSPA's watersheds. The Authority has initiated a stream baseflow assessment program. The main objective is to obtain baseflow information to help develop a long-term baseflow monitoring network using a predetermined distribution of measurement sites. These data are also necessary for model calibration in water budgeting exercises, a necessary component for SWP activities.

Table B-7 lists sites where streamflow measurements have been taken annually, beginning in 2002. The field program measures flow taken over spring/summer/fall seasons. Field flow measurements are generally taken at stream crossings and stream gauge stations. These measurements represent a significant source of information that supports aquatic studies, groundwater discharge, and water budgets, including numerical model calibration. Linking low-flow measurements to streamflow gauges was undertaken by CLOSPA between 2006 and 2007 as part of SWP activities.

| SITE | LOT | CON | WATER-SHED | SITE | LOT | CON | WATER-SHED | SITE | LOT | CON | WATER-SHED |
|-------|-----|-------|-------------------|-------|-----|-------|----------------|-------|-----|-----|--------------|
| SBF1 | 10 | B.F.C | Soper Creek | FBF4 | 28 | 06 | Farewell Creek | OBF34 | 15 | 09 | Oshawa Creek |
| SBF2 | 07 | 01 | | FBF5 | 30 | 06 | | OBF35 | 18 | 07 | |
| SBF3 | 04 | 01 | | FBF6 | 29 | 07 | | OBF36 | 17 | 08 | |
| SBF5 | 08 | 02 | | HBF1 | 04 | 01 | Harmony Creek | OBF37 | 19 | 07 | |
| SBF6 | 06 | 03 | | HBF2 | 05 | 02 | | OBF38 | 19 | 09 | |
| SBF7 | 05 | 05 | | HBF3 | 06 | 02 | | LBF1 | 30 | 01 | |
| SBF8 | 07 | 05 | | HBF4 | 34 | 04 | | LBF2 | 28 | 03 | |
| SBF9 | 07 | 07 | | HBF5 | 32 | 04 | | LBF3 | 26 | 04 | |
| SBF10 | 07 | 08 | | HBF6 | 35 | 04 | | LBF4 | 23 | 06 | |
| SBF11 | 07 | 08 | | HBF7 | 03 | 03 | | LBF5 | 22 | 07 | |
| SBF12 | 03 | 05 | | HBF8 | 03 | 04 | | LBF6 | 21 | 08 | |
| SBF13 | 02 | 06 | | HBF9 | 01 | 04 | | LBF7 | 24 | 07 | |
| SBF14 | 03 | 06 | | HBF10 | 34 | 05 | | LBF8 | 24 | 08 | |
| SBF15 | 03 | 06 | | OBF1 | 13 | 04 | LBF9 | 24 | 09 | | |
| SBF16 | 03 | 07 | | OBF2 | 11 | 04 | LBF10 | 24 | 09 | | |
| SBF17 | 05 | 07 | OBF3 | 06 | 06 | LBF11 | 25 | 09 | | | |
| BBF1 | 14 | 02 | Bowmanville Creek | OBF4 | 06 | 06 | Oshawa Creek | LBF12 | 25 | 08 | Lynde Creek |
| BBF2 | 15 | 04 | | OBF5 | 03 | 06 | | LBF13 | 27 | 09 | |
| BBF3 | 12 | 05 | | OBF6 | 35 | 08 | | LBF14 | 30 | 09 | |
| BBF4 | 11 | 07 | | OBF7 | 32 | 08 | | LBF15 | 04 | 01 | |
| BBF5 | 10 | 08 | | OBF8 | 09 | 06 | | LBF16 | 04 | 01 | |
| BBF6 | 14 | 05 | | OBF9 | 06 | 06 | | LBF17 | 32 | 03 | |
| BBF7 | 14 | 06 | | OBF10 | 06 | 07 | | LBF18 | 31 | 05 | |
| BBF8 | 12 | 08 | | OBF11 | 08 | 06 | | LBF19 | 28 | 06 | |
| BBF9 | 15 | 06 | | OBF12 | 08 | 08 | | LBF20 | 27 | 07 | |
| BBF10 | 14 | 08 | | OBF13 | 02 | 09 | | LBF21 | 29 | 06 | |
| BBF11 | 14 | 09 | | OBF14 | 03 | 08 | | LBF22 | 30 | 07 | |
| BBF12 | 15 | 08 | | OBF15 | 06 | 09 | | LBF23 | 31 | 08 | |
| BBF13 | 16 | 05 | | OBF16 | 07 | 08 | | LBF24 | 32 | 09 | |
| BBF14 | 17 | 06 | | OBF17 | 10 | 06 | | LBF25 | 32 | 06 | |
| BBF15 | 18 | 05 | | OBF18 | 14 | 04 | | LBF26 | 31 | 07 | |
| BBF16 | 19 | 06 | | OBF19 | 13 | 05 | | LBF27 | 34 | 08 | |
| BBF17 | 20 | 06 | | OBF20 | 18 | 05 | | LBF28 | 34 | 08 | |
| BBF18 | 20 | 07 | | OBF21 | 13 | 07 | | LBF29 | 03 | 07 | |
| BBF19 | 21 | 08 | | OBF22 | 12 | 07 | | LBF30 | 02 | 08 | |
| BBF20 | 28 | 08 | | OBF23 | 10 | 08 | | LBF31 | 32 | 06 | |
| BBF21 | 26 | 09 | | OBF24 | 11 | 08 | | LBF32 | 35 | 06 | |
| BBF22 | 16 | 05 | | OBF25 | 12 | 08 | | LBF33 | 31 | 01 | |
| BLBF1 | 24 | 03 | Black Creek | OBF26 | 09 | 08 | LBF34 | 34 | 01 | | |
| BLBF2 | 20 | 04 | | OBF27 | 11 | 08 | LBF35 | 34 | 03 | | |
| BLBF3 | 21 | 04 | | OBF28 | 10 | 08 | LBF36 | 34 | 05 | | |
| BLBF4 | 24 | 05 | | OBF29 | 15 | 07 | LBF37 | 01 | 03 | | |
| BLBF5 | 24 | 06 | | OBF30 | 14 | 08 | LBF38 | 02 | 05 | | |
| FBF1 | 30 | 04 | Farewell Creek | OBF31 | 14 | 08 | LBF39 | 03 | 05 | | |
| FBF2 | 26 | 05 | | OBF32 | 14 | 09 | | | | | |
| FBF3 | 25 | 06 | | OBF33 | 15 | 08 | | | | | |

Table B-7: Low-Flow Monitoring Details, CLOSPA Stations

Additionally, a low-flow survey was conducted under the ORMGP initiative in 2002 by Conestoga Rovers and Associates. Data gathered from the survey, which includes 47 locations within CLOSPA's jurisdiction, are stored in the ORMGP database.

B1.2.7 Biological Monitoring

Biological sampling measures ecological effects, whereas sampling for chemical and physical parameters measures stressors (i.e., environmental contamination). Though source water protection technical guidelines do not directly link the assessment and protection of drinking water to biological assessment, it is recognized that the various components of the watershed are closely linked. Protecting source water is important to the biological health of the watershed, and biological indicators are fundamental in protecting source water. CLOSPA's biological surveys involve sampling creatures, such as benthic macroinvertebrates and fish, found living within the aquatic environment. Benthic macroinvertebrates make good health indicators of aquatic ecosystems because:

- They generally have limited mobility, which makes them vulnerable to many creek stresses that may occur;
- They have short life cycles;
- They are easily collected and identified; and
- Their spatial distribution across the watershed is good.

Historically, CLOSPA's aquatic biological sampling has followed the BioMap protocol (Griffiths, 1999). Sampling was undertaken as part of the Aquatic Resource Management Plan (ARMP) activities for all watersheds within the study area. In CLOSPA, ARMPs have been developed on a watershed-by-watershed basis. To date, two ARMPs have been completed for CLOSPA with three additional ARMPs in progress for a total of five to cover the full CLOSPA jurisdiction.

To coordinate long-term monitoring efforts, CLOSPA joined the Ontario Benthos Biomonitoring Network (OBBN) in 2005. OBBN sites (**Figure B-2**) are closely linked to CLOSPA's PWQMN sites. This provincial network allows CLOSPA to follow a standardized methodology, share resources, and offer technical support.

B1.2.8 Coastal Wetland Monitoring

The Durham Region Coastal Wetland Monitoring Project is designed as a long-term program that will assess the health of 15 wetlands along the north shore of Lake Ontario in the Durham Region (**Figure B-3** and **Table B-8**).

To standardize the collection of biological and physical data among the partner organizations, a methodology handbook was developed by Environment Canada and the Central Lake Ontario Conservation Authority and fieldwork began in the spring of 2002.

Water levels in the Great Lakes have been recorded by the Canadian Hydrographic Service since 1860. These data show that levels in Lake Ontario have varied by up to two metres since that time. In 1958, however, lake level regulation was implemented, which moderated levels. While lake levels still fluctuate, they do not do so to the extent that occurred prior to regulation.




Figure B-2: Ontario Benthic Biomonitoring Network (OBBN) within the Study Area




Figure B-3: Coastal Wetland Monitoring Project Locations, Durham Region

| # | Wetland | SPA | # | Wetland | SPA* |
|---|------------------------|--------|---|------------------------|--------|
| 1 | Rouge River Marsh | TRSPA | 9 | Pumphouse Marsh | CLOSPA |
| 2 | Frenchman's Bay Marsh | TRSPA | 10 | Oshawa Second Marsh | CLOSPA |
| 3 | Hydro Marsh | TRSPA | 11 | McLaughlin Bay Marsh | CLOSPA |
| 4 | Duffins Creek Marsh | TRSPA | 12 | Westside Marsh | CLOSPA |
| 5 | Carruthers Creek Marsh | TRSPA | 13 | Bowmanville Marsh | CLOSPA |
| 6 | Cranberry Marsh | CLOSPA | 14 | Wilmot Creek Marsh | GRSPA |
| 7 | Lynde Creek Marsh | CLOSPA | 15 | Port Newcastle Wetland | GRSPA |
| 8 | Corbett Creek Marsh | CLOSPA | *TRCA: Toronto and Region Source Protection Area CLOSPA: Central Lake Ontario Source Protection Area GRSPA: Ganaraska Region Source Protection Area | | |

Table B-8: Durham Region Coastal Wetlands Currently Monitored

The Durham Region Coastal Wetlands Project monitors both physical features and biological communities. The following physical features or aspects are observed within the Coastal Wetland Monitoring Program:

- **Water Quality**—Measure various water quality parameters, including turbidity (clarity of water), conductivity, nitrogen, and phosphorus;
- **Water Levels**—For wetlands that can be cut off from Lake Ontario due to the formation of a barrier beach, measure water levels throughout the vegetation growing season (May to October);
- **Sediment Quality**—Collect recently deposited sediments to analyze for various contaminants including pesticides, metals, PCBs and PAHs;
- **Bathymetry**—Map wetland basin topography to reveal contours;
- **Watershed Vegetation**—Ecological Land Classification to Community Series level summarized for each wetland's watershed;
- **Land-use Change in Adjacent Uplands**—Compare current land use in 1000 meter zone around wetland with expected land use according to municipal and regional Official Plans. Obtain percentages of change for each land use category; and
- **Land-use Change in Watershed**—In conjunction with Watershed Management Plans, compare current land usage with expected land usage according to municipal and regional Official Plans Sediment and Nutrient Loading Computer modelling incorporating a Digital Elevation Model (DEM). This step is to be completed when technology becomes available.

The following biological communities are observed within the Coastal Wetland Monitoring Program:

- **Birds**—Survey marsh breeding bird communities using the Marsh Monitoring Program methodology;
- **Amphibians**—Survey amphibian communities using the Marsh Monitoring Program methodology;
- **Fish**—Survey wetland fish community using electrofishing boat;
- **Macroinvertebrates**—Sample wetland macroinvertebrates by sweep-netting through water column;
- **Wetland Vegetation**—Use Ecological Land Classification to define vegetation communities at each wetland and surrounding 500 metres;
- **Submerged Plants**—Sample submerged aquatic vegetation using 20 randomly placed quadrants; and
- **Identifying Key Habitats**—Over time, identify and track habitats associated with species at risk (i.e., endangered, threatened, or of special concern).

B1.3 INFORMATION MANAGEMENT SYSTEM

Sources of information (details and descriptions) identified through source protection activities are to be tracked through CLOSPA's current information management system (IMS). IMS is an updatable and searchable database that contains metadata related to reports, documents, and correspondence. Folders are assigned IMS numbers (ID #) and updated information related to a particular folder is linked using the same IMS number as the parent folder.

The database is searchable by keyword, municipality, watershed, name, address, municipality number, permit number, date/owner, and folder ID number or attachment ID number. Information added to the database may contain a description of the report or data. Folders may be linked or cross-referenced by ID numbers. CLOSPA's IMS is regularly backed up and the database is accessible through the LAN by an IMS interface loaded on each workstation. Centralized updates or edits to IMS are typically required to maintain standardization of format within the system. IMS directly links digital files through the local area network. Hard copies of information are tracked in IMS to facilitate accessibility either within individual office cabinets or in the main administrative file location.

B1.4 METHODS OF ANALYSIS

The Watershed Characterization is a description of the local watershed area and was developed by compiling all the available information about the area. It will include topics such as watershed features, the water quality, the wells and intakes that draw drinking water, and the natural and human-made influences. Maps were produced to provide a visualization of the watershed. This information gathering process will be iterative and continuous and will occur wherever possible to enhance the available data.

The watershed features include topography, physiography, geology, hydrology (surface water flow system) and hydrogeology (groundwater flow system), ecology, naturally vegetated areas, and climate. This information provides the background necessary for a more in-depth analysis in subsequent phases of the Assessment Report, including the Water Budget and Stress Assessment, the Vulnerability Analysis, and the Summary of Threats and Issues.

The water quality conditions and long-term trends in the watershed were identified. Maps and graphics are used to illustrate these trends. The objective was to describe the quality of surface water and groundwater using existing information and to determine whether the water quality is improving, deteriorating, or remaining constant.

The current water use was inventoried, as were historical takings, to illustrate where most of the water is going and at what times during the year. The inventory estimated population growth in the watershed area, which has a significant impact on future water demands.

The SPA also identified land-use activities that are known to pose a threat to the quality or quantity of drinking water to determine human and ecological impacts.

A Watershed Characterization Report has been prepared for the Source Protection Areas (Central Lake Ontario Conservation Authority, 2007). Workshops involving the conservation authority and municipal partners were held in late 2006 and early 2007 to review the contents of earlier versions of these reports. The most recent versions include edits and updates that are the result of the comments provided. The Province has established a panel to review the Watershed Characterization Reports. Comments from this review panel were received on January 8, 2008, and were incorporated into the final Assessment Report.

B1.5 SURFACE WATER QUALITY DATA ANALYSIS AND REPORTING

The analysis and reporting of surface water quality data were accomplished in three steps:

- Exploratory analysis;
- Statistical analysis; and
- Reporting results.

B1.5.1 Exploratory Analysis

The first step involves plotting water quality observations to visually examine the attributes of the data (e.g., outliers and data entry errors). Each water quality observation is represented as a single point or dot. The y-axis (the dependent axis) is the concentration of a water quality parameter, and the x-axis (the independent axis) is time, usually represented as months or years. Specifically, a plot of water quality results against time allows for the:

- Observation of seasonal and annual trends;
- Identification of anomalous results and potential errors;
- Comparison of results to water quality criteria (e.g., Provincial Water Quality Objectives, Canadian Water Quality Guidelines);
- Observation of changes in water quality over time;
- Identification of missing periods of record (data gaps); and
- Identification of biases introduced by the timing of water quality measurements.

B1.5.2 Statistical Analysis

The second step in the analysis of surface water quality data involves the selection and application of statistical tests to establish the significance of differences, trends, and relationships that were identified in the exploration of the data.

B1.5.3 Reporting Results

The third step involves the use of graphics such as maps and boxplots to present selected results in a format consistent with the information needs and technical knowledge of the target audience. Results that are selected for reporting should describe the prevailing surface water quality conditions in the watershed (**Table B-9**).

| Parameter | Provincial Limits | Unit | Number of Samples | Minimum | Maximum | Arithmetic Mean | Standard Deviation | 10th Percentile | Median 50th Percentile | 90th Percentile | Mann-Kendall Statistic indicating increasing or decreasing trend | Mann-Kendall Statistic, approximate Z value for calculating probability | Mann-Kendall Result 95% significance |
|--------------------------------------|-------------------|------|-------------------|---------|---------|-----------------|--------------------|-----------------|------------------------|-----------------|--|---|--------------------------------------|
| SWQ1 (Lynde) | | | | | | | | | | | | | |
| Phosphorus, Total | 30 | µg/l | 69 | 8 | 280 | 39.80 | 37 | 17.800 | 31 | 68.40 | 7 | 0.031 | No Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 23 | 0.084 | 1.59 | 0.63 | 0.352 | 0.229 | 0.58 | 1.08 | 83 | 2.170 | Increasing Trend |
| Nitrate as N | 10 | mg/l | 47 | 0.17 | 2.64 | 0.79 | 0.526 | 0.277 | 0.62 | 1.49 | 77 | 0.700 | No Trend |
| Cl | 250 | mg/l | 70 | 26 | 661 | 97.20 | 79.400 | 56.200 | 86.80 | 116.40 | 179 | 0.900 | No Trend |
| Cu | 5 | µg/l | 70 | 0.200 | 5.80 | 1.53 | 0.960 | 0.630 | 1.33 | 2.46 | 174 | 0.880 | No Trend |
| Cd | 0.1 - 0.5 | µg/l | 53 | 0.100 | 1.09 | 0.33 | 0.337 | 0.100 | 0.10 | 0.88 | 164 | 1.250 | No Trend |
| Co | 0.9 | µg/l | 62 | 0.007 | 1.47 | 0.42 | 0.332 | 0.200 | 0.30 | 0.81 | 278 | 1.680 | No Trend |
| Fe | 0.3 | mg/l | 70 | 0.003 | 1.17 | 0.26 | 0.204 | 0.104 | 0.21 | 0.37 | 164 | 0.830 | No Trend |
| Ni | 25 | µg/l | 63 | 0.064 | 3.96 | 0.85 | 0.730 | 0.100 | 0.83 | 1.38 | 116 | 0.680 | No Trend |
| Pb | 5 - 25 | µg/l | 55 | 0.043 | 6 | 1.32 | 1.440 | 0.140 | 0.70 | 3.23 | 285 | 2.060 | Decreasing Trend |
| Zn | 30 | µg/l | 69 | 0.100 | 18 | 2.87 | 3.040 | 0.660 | 2.10 | 5.20 | 45 | 0.228 | No Trend |
| SWQ8 (Lynde) | | | | | | | | | | | | | |
| Phosphorus, Total | 30 | µg/l | 119 | 3 | 562 | 37 | 74.700 | 6 | 18 | 66 | 770 | 1.770 | No Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 25 | 0.554 | 1.21 | 0.774 | 0.158 | 0.610 | 0.746 | 0.95 | 40 | 0.91 | No Trend |
| Nitrate as N | 10 | mg/l | 46 | 0.455 | 1.78 | 0.910 | 0.340 | 0.614 | 0.815 | 1.43 | 126 | 1.18 | No Trend |
| Cl | 250 | mg/l | 119 | 9.400 | 209 | 37.600 | 24.400 | 13.600 | 39.400 | 56.90 | 3854 | 8.85 | Increasing Trend |
| Cu | 5 | µg/l | 93 | 0.089 | 7 | 0.940 | 1.080 | 0.200 | 0.600 | 1.98 | 438 | 1.45 | No Trend |
| Cd | 0.1 - 0.5 | µg/l | 46 | 0.075 | 1.14 | 0.208 | 0.226 | 0.100 | 0.100 | 0.50 | 337 | 3.18 | Decreasing Trend |
| Co | 0.9 | µg/l | 45 | 0.038 | 2.4 | 0.350 | 0.353 | 0.200 | 0.200 | 0.56 | 59 | 0.57 | No Trend |
| Fe | 0.3 | mg/l | 96 | 0.001 | 4.36 | 0.211 | 0.530 | 0.054 | 0.092 | 0.37 | 579 | 1.83 | No Trend |
| Ni | 25 | µg/l | 46 | 0.100 | 6 | 0.650 | 0.980 | 0.100 | 0.400 | 1 | 66 | 0.62 | No Trend |
| Pb | 5 - 25 | µg/l | 86 | 0.005 | 13 | 3.230 | 2.780 | 0.100 | 5 | 5 | 1870 | 7 | Decreasing Trend |
| Zn | 30 | µg/l | 88 | 0.018 | 220 | 6.340 | 23.800 | 0.395 | 1.750 | 12.90 | 1279 | 4.60 | No Trend |

| SWQ9 (Lynde) | | | | | | | | | | | | | |
|--------------------------------------|-----------|------|----|--------|-------|--------|--------|--------|--------|-------|------|-------|------------------|
| Phosphorus, Total | 30 | µg/l | 43 | 6 | 470 | 35.700 | 69.2 | 13 | 22 | 41.60 | -78 | -0.81 | No Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 0 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | 0 | 0 | No Trend |
| Nitrate as N | 10 | mg/l | 43 | 0.280 | 3.03 | 1.030 | 0.510 | 0.596 | 0.860 | 1.63 | -25 | -0.25 | No Trend |
| Cl | 250 | mg/l | 43 | 19.700 | 131 | 48.200 | 20.300 | 32.500 | 42.100 | 57.90 | 100 | 1.04 | No Trend |
| Cu | 5 | µg/l | 43 | 0.200 | 8 | 0.720 | 1.170 | 0.200 | 0.600 | 1 | 200 | 2.08 | Increasing Trend |
| Cd | 0.1 - 0.5 | µg/l | 43 | 0.100 | 0.50 | 0.112 | 0.0625 | 0.100 | 0.100 | 0.10 | -35 | -0.36 | No Trend |
| Co | 0.9 | µg/l | 43 | 0.200 | 1.90 | 0.274 | 0.275 | 0.200 | 0.200 | 0.30 | 256 | 2.67 | No Trend |
| Fe | 0.3 | mg/l | 43 | 0.0333 | 3.18 | 0.207 | 0.466 | 0.088 | 0.133 | 0.20 | -156 | -1.62 | No Trend |
| Ni | 25 | µg/l | 43 | 0.100 | 4 | 0.460 | 0.71 | 0.10 | 0.20 | 0.96 | 456 | 4.76 | Increasing Trend |
| Pb | 5 - 25 | µg/l | 43 | 0.100 | 8 | 0.680 | 1.10 | 0.10 | 0.70 | 0.70 | -369 | -3.85 | Decreasing Trend |
| Zn | 30 | µg/l | 43 | 0.100 | 20 | 1.200 | 3.03 | 0.20 | 0.50 | 1.66 | -92 | -0.95 | No Trend |
| SWQ2 (Oshawa) | | | | | | | | | | | | | |
| Phosphorus, Total | 30 | µg/l | 68 | 6 | 115 | 26.100 | 21.40 | 8.70 | 18.50 | 47 | 67 | 0.35 | No Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 23 | 0.397 | 2.30 | 0.840 | 0.37 | 0.52 | 0.82 | 1.09 | 65 | 1.69 | Increasing Trend |
| Nitrate as N | 10 | mg/l | 46 | 0.590 | 3.04 | 1.090 | 0.55 | 0.66 | 0.84 | 1.79 | 89 | 0.83 | No Trend |
| Cl | 250 | mg/l | 69 | 10.800 | 200 | 68.800 | 29.70 | 45 | 62.10 | 89.30 | 351 | 1.81 | Increasing Trend |
| Cu | 5 | µg/l | 69 | 0.200 | 9.40 | 1.430 | 1.38 | 0.30 | 1 | 2.52 | 310 | 1.60 | No Trend |
| Cd | 0.1 - 0.5 | µg/l | 58 | 0.029 | 1.54 | 0.328 | 0.35 | 0.10 | 0.10 | 0.82 | 156 | 1.04 | No Trend |
| Co | 0.9 | µg/l | 57 | 0.237 | 1.77 | 0.505 | 0.44 | 0.20 | 0.30 | 1.19 | 60 | 0.41 | No Trend |
| Fe | 0.3 | mg/l | 69 | 0.001 | 0.57 | 0.193 | 0.12 | 0.09 | 0.18 | 0.36 | 80 | 0.41 | No Trend |
| Ni | 25 | µg/l | 68 | 0.081 | 25.50 | 1.820 | 3.38 | 0.37 | 1.25 | 2.20 | 377 | 2 | Increasing Trend |
| Pb | 5 - 25 | µg/l | 51 | 0.099 | 11.20 | 1.590 | 2.16 | 0.10 | 0.70 | 3.50 | 33 | 0.26 | No Trend |
| Zn | 30 | µg/l | 65 | 0.400 | 34.60 | 3.400 | 4.76 | 0.81 | 1.98 | 6.70 | 256 | 1.44 | No Trend |
| SWQ10 (Oshawa) | | | | | | | | | | | | | |
| Phosphorus, Total | 30 | µg/l | 68 | 5 | 130 | 17.600 | 20.30 | 6 | 11.50 | 33.20 | 18 | 0.09 | No Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 23 | 0.524 | 1.27 | 0.760 | 0.190 | 0.565 | 0.73 | 0.98 | 65 | 1.69 | Increasing Trend |
| Nitrate as N | 10 | mg/l | 45 | 0.370 | 2.26 | 0.950 | 0.400 | 0.650 | 0.76 | 1.54 | 176 | 1.71 | Decreasing Trend |
| Cl | 250 | mg/l | 68 | 22.400 | 80 | 33.100 | 9.700 | 25.100 | 29.80 | 43.70 | 513 | 2.70 | Increasing Trend |
| Cu | 5 | µg/l | 50 | 0.057 | 5 | 0.570 | 0.730 | 0.200 | 0.40 | 0.90 | 32 | 0.26 | No Trend |

| | | | | | | | | | | | | | |
|--------------------------------------|-----------|------|----|--------|------|--------|--------|-------|-------|--------|------|-------|------------------|
| Cd | 0.1 - 0.5 | µg/l | 44 | 0.012 | 0.94 | 0.142 | 0.147 | 0.100 | 0.10 | 0.21 | 67 | 0.67 | No Trend |
| Co | 0.9 | µg/l | 43 | 0.024 | 1.4 | 0.305 | 0.272 | 0.169 | 0.20 | 0.50 | 6 | 0.05 | No Trend |
| Fe | 0.3 | mg/l | 54 | 0.056 | 0.64 | 0.127 | 0.103 | 0.068 | 0.10 | 0.18 | 191 | 1.42 | No Trend |
| Ni | 25 | µg/l | 48 | 0.100 | 2.13 | 0.445 | 0.490 | 0.100 | 0.20 | 0.93 | 105 | 0.92 | No Trend |
| Pb | 5 - 25 | µg/l | 45 | 0.100 | 10 | 1.540 | 2.360 | 0.100 | 0.70 | 3.90 | 495 | 4.83 | Decreasing Trend |
| Zn | 30 | µg/l | 43 | 0.057 | 11.2 | 0.990 | 1.880 | 0.178 | 0.40 | 1.84 | 16 | 0.16 | No Trend |
| SWQ11 (Oshawa) | | | | | | | | | | | | | |
| Phosphorus, Total | 30 | µg/l | 42 | 6 | 230 | 16.200 | 35.100 | 6 | 6 | 30.30 | 12 | 0.12 | No Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 0 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | 0 | 0 | No Trend |
| Nitrate as N | 10 | mg/l | 42 | 0.760 | 2.06 | 1.150 | 0.282 | 0.910 | 1.07 | 1.56 | -256 | -2.76 | Decreasing Trend |
| Cl | 250 | mg/l | 42 | 21.100 | 38 | 25.100 | 3.370 | 22 | 23.80 | 29.50 | 193 | 2.08 | Increasing Trend |
| Cu | 5 | µg/l | 42 | 0.200 | 5 | 0.550 | 0.756 | 0.200 | 0.40 | 0.89 | 181 | 1.95 | Increasing Trend |
| Cd | 0.1 - 0.5 | µg/l | 42 | 0.100 | 1 | 0.133 | 0.151 | 0.100 | 0.10 | 0.10 | -6 | -0.05 | No Trend |
| Co | 0.9 | µg/l | 42 | 0.10 | 1.60 | 0.28 | 0.257 | 0.20 | 0.20 | 0.30 | 140 | 1.51 | No Trend |
| Fe | 0.3 | mg/l | 42 | 0.06 | 1.43 | 0.13 | 0.214 | 0.06 | 0.08 | 0.20 | -205 | -2.20 | Decreasing Trend |
| Ni | 25 | µg/l | 42 | 0.10 | 2 | 0.34 | 0.46 | 0.10 | 0.10 | 0.79 | 397 | 4.30 | Increasing Trend |
| Pb | 5 - 25 | µg/l | 42 | 0.10 | 1.10 | 0.56 | 0.284 | 0.10 | 0.70 | 0.70 | -291 | -3.15 | Decreasing Trend |
| Zn | 30 | µg/l | 42 | 0.10 | 6.50 | 0.82 | 1.35 | 0.20 | 0.40 | 1.93 | -7 | -0.07 | No Trend |
| SWQ12 (Harmony) | | | | | | | | | | | | | |
| Phosphorus, Total | 30 | µg/l | 9 | 8 | 66 | 28.10 | 19.4 | 8 | 25 | 50 | 5 | 0.42 | No Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 0 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | 0 | 0 | No Trend |
| Nitrate as N | 10 | mg/l | 9 | 0.75 | 2.28 | 1.17 | 0.484 | 0.79 | 1 | 1.66 | -2 | -0.10 | No Trend |
| Cl | 250 | mg/l | 9 | 42 | 413 | 228.90 | 121.5 | 91.60 | 244 | 363.40 | -14 | -1.36 | No Trend |
| Cu | 5 | µg/l | 9 | 1 | 4.10 | 2.06 | 0.88 | 1.32 | 1.90 | 2.90 | 5 | 0.42 | No Trend |
| Cd | 0.1 - 0.5 | µg/l | 9 | 0.10 | 0.10 | 0.10 | 0 | 0.10 | 0.10 | 0.10 | 0 | 0 | No Trend |
| Co | 0.9 | µg/l | 9 | 0.20 | 0.50 | 0.29 | 0.105 | 0.20 | 0.30 | 0.42 | 15 | 1.46 | No Trend |
| Fe | 0.3 | mg/l | 9 | 0.14 | 0.35 | 0.21 | 0.071 | 0.15 | 0.19 | 0.30 | 2 | 0.10 | No Trend |
| Ni | 25 | µg/l | 9 | 0.10 | 1.70 | 0.86 | 0.53 | 0.10 | 0.90 | 1.38 | 23 | 2.30 | Increasing Trend |
| Pb | 5 - 25 | µg/l | 9 | 0.20 | 0.70 | 0.56 | 0.219 | 0.28 | 0.70 | 0.70 | -16 | -1.60 | No Trend |
| Zn | 30 | µg/l | 9 | 1.30 | 7 | 3.30 | 1.97 | 1.54 | 2.30 | 5.90 | 15 | 1.46 | No Trend |

| SWQ3 (Farewell) | | | | | | | | | | | | | |
|--------------------------------------|-----------|------|----|--------|-------|--------|--------|-------|-------|--------|------|-------|------------------|
| Phosphorus, Total | 30 | µg/l | 68 | 6 | 1310 | 59.50 | 160.70 | 12 | 28 | 81.40 | 223 | 1.18 | Decreasing Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 22 | 0.650 | 3.40 | 1.41 | 0.57 | 0.74 | 1.35 | 1.81 | 4 | 0.09 | No Trend |
| Nitrate as N | 10 | mg/l | 46 | 0.750 | 5.23 | 1.57 | 0.78 | 1 | 1.34 | 2.54 | 231 | 2.18 | Decreasing Trend |
| Cl | 250 | mg/l | 68 | 32.400 | 594 | 142.90 | 77.30 | 82.60 | 134 | 191.60 | 7 | 0.03 | Increasing Trend |
| Cu | 5 | µg/l | 68 | 0.200 | 67.80 | 2.90 | 8.10 | 0.70 | 1.65 | 3.76 | 241 | 1.27 | Decreasing Trend |
| Cd | 0.1 - 0.5 | µg/l | 56 | 0.010 | 1.51 | 0.30 | 0.35 | 0.10 | 0.10 | 0.72 | 288 | 2.03 | No Trend |
| Co | 0.9 | µg/l | 51 | 0.120 | 2 | 0.46 | 0.38 | 0.20 | 0.30 | 0.91 | 25 | 0.20 | No Trend |
| Fe | 0.3 | mg/l | 68 | 0.001 | 2.04 | 0.26 | 0.27 | 0.13 | 0.21 | 0.36 | 116 | 0.61 | Decreasing Trend |
| Ni | 25 | µg/l | 66 | 0.0001 | 50 | 1.50 | 6.10 | 0.10 | 0.72 | 1.30 | 174 | 0.96 | No Trend |
| Pb | 5 - 25 | µg/l | 51 | 0.027 | 67.30 | 3.04 | 9.60 | 0.20 | 0.70 | 4.75 | 423 | 3.43 | Decreasing Trend |
| Zn | 30 | µg/l | 68 | 0.250 | 394 | 9.40 | 47.60 | 0.95 | 2.20 | 7 | 410 | 2.17 | Decreasing Trend |
| SWQ13 (Farewell) | | | | | | | | | | | | | |
| Phosphorus, Total | 30 | µg/l | 42 | 6 | 150 | 19.10 | 24.50 | 6.10 | 12 | 30.70 | 124 | 1.33 | No Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 0 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | 0 | 0 | No Trend |
| Nitrate as N | 10 | mg/l | 42 | 0.740 | 4.80 | 1.86 | 0.59 | 1.44 | 1.77 | 2.40 | -102 | -1.10 | No Trend |
| Cl | 250 | mg/l | 42 | 18.20 | 63.30 | 29.60 | 10 | 20.70 | 27 | 38.70 | 96 | 1.03 | No Trend |
| Cu | 5 | µg/l | 42 | 0.200 | 3 | 0.676 | 0.560 | 0.20 | 0.55 | 1.10 | 201 | 2.17 | Increasing Trend |
| Cd | 0.1 - 0.5 | µg/l | 42 | 0.100 | 0.40 | 0.107 | 0.046 | 0.10 | 0.10 | 0.10 | -25 | -0.26 | No Trend |
| Co | 0.9 | µg/l | 42 | 0.100 | 0.50 | 0.233 | 0.082 | 0.20 | 0.20 | 0.30 | 237 | 2.56 | No Trend |
| Fe | 0.3 | mg/l | 42 | 0.033 | 0.49 | 0.089 | 0.078 | 0.04 | 0.07 | 0.15 | 18 | 0.18 | No Trend |
| Ni | 25 | µg/l | 42 | 0.100 | 1.20 | 0.255 | 0.253 | 0.10 | 0.10 | 0.49 | 410 | 4.43 | Increasing Trend |
| Pb | 5 - 25 | µg/l | 42 | 0.100 | 0.90 | 0.543 | 0.259 | 0.10 | 0.70 | 0.70 | -337 | -3.65 | Decreasing Trend |
| Zn | 30 | µg/l | 42 | 0.100 | 18.30 | 1.29 | 2.900 | 0.20 | 0.50 | 1.79 | 281 | 3.04 | Increasing Trend |
| SWQ14 (Black) | | | | | | | | | | | | | |
| Phosphorus, Total | 30 | µg/l | 67 | 6 | 318 | 73.700 | 55.500 | 21 | 64 | 143.20 | 70 | 0.37 | No Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 21 | 0.410 | 1.75 | 0.905 | 0.323 | 0.67 | 0.84 | 1.29 | 66 | 1.96 | Decreasing Trend |
| Nitrate as N | 10 | mg/l | 46 | 0.200 | 3.28 | 1 | 0.540 | 0.50 | 0.82 | 1.64 | 247 | 2.33 | Decreasing Trend |
| Cl | 250 | mg/l | 67 | 21.600 | 151 | 72.600 | 30.400 | 40.40 | 63.40 | 115 | 99 | 0.53 | No Trend |
| Cu | 5 | µg/l | 54 | 0.109 | 6.30 | 1.150 | 1.160 | 0.21 | 0.87 | 2.30 | 308 | 2.3 | Increasing Trend |

**Approved Assessment Report:
Central Lake Ontario Source Protection Area**

**Appendix B1:
Watershed Characterization**

| | | | | | | | | | | | | | |
|--------------------------------------|-----------|------|----|---------|-------|--------|--------|-------|--------|-------|-----|------|-------------------------|
| Cd | 0.1 - 0.5 | µg/l | 44 | 0.030 | 1.86 | 0.200 | 0.307 | 0.10 | 0.10 | 0.48 | 200 | 2.01 | Decreasing Trend |
| Co | 0.9 | µg/l | 44 | 0.0187 | 1.59 | 0.307 | 0.260 | 0.20 | 0.20 | 0.52 | 9 | 0.08 | No Trend |
| Fe | 0.3 | mg/l | 54 | 0.051 | 0.58 | 0.184 | 0.103 | 0.10 | 0.16 | 0.32 | 70 | 0.52 | No Trend |
| Ni | 25 | µg/l | 49 | 0.014 | 1.20 | 0.510 | 0.345 | 0.10 | 0.50 | 0.97 | 131 | 1.12 | No Trend |
| Pb | 5 - 25 | µg/l | 45 | 0.0710 | 15 | 1.42 | 2.50 | 0.10 | 0.700 | 3.32 | 332 | 3.24 | Decreasing Trend |
| Zn | 30 | µg/l | 54 | 0.3600 | 20.10 | 4.33 | 3.90 | 1.53 | 2.950 | 9.70 | 96 | 0.71 | No Trend |
| SWQ4 (Bowmanville) | | | | | | | | | | | | | |
| Phosphorus, Total | 30 | µg/l | 65 | 6 | 165 | 23.80 | 25.20 | 6 | 15 | 60 | 91 | 0.51 | Decreasing Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 19 | 0.3930 | 1.93 | 0.79 | 0.43 | 0.41 | 0.636 | 1.46 | 39 | 1.33 | No Trend |
| Nitrate as N | 10 | mg/l | 46 | 0.4000 | 3.97 | 0.99 | 0.68 | 0.49 | 0.660 | 1.75 | 112 | 1.05 | No Trend |
| Cl | 250 | mg/l | 65 | 13.6000 | 282 | 28.70 | 33 | 16.30 | 23.500 | 36.10 | 480 | 2.70 | Increasing Trend |
| Cu | 5 | µg/l | 65 | 0.0262 | 2.1 | 0.74 | 0.44 | 0.20 | 0.700 | 1.34 | 614 | 3.47 | Decreasing Trend |
| Cd | 0.1 - 0.5 | µg/l | 58 | 0.0010 | 1.31 | 0.28 | 0.32 | 0.10 | 0.100 | 0.82 | 29 | 0.19 | Decreasing Trend |
| Co | 0.9 | µg/l | 53 | 0.0860 | 1.71 | 0.36 | 0.32 | 0.20 | 0.200 | 0.66 | 76 | 0.58 | No Trend |
| Fe | 0.3 | mg/l | 66 | 0.0005 | 0.44 | 0.13 | 0.10 | 0.06 | 0.106 | 0.23 | 87 | 0.48 | Decreasing Trend |
| Ni | 25 | µg/l | 57 | 0.0292 | 2.12 | 0.37 | 0.44 | 0.10 | 0.191 | 0.97 | 261 | 1.79 | Decreasing Trend |
| Pb | 5 - 25 | µg/l | 51 | 0.0272 | 21.4 | 2.30 | 4.24 | 0.10 | 0.700 | 5.17 | 338 | 2.74 | Decreasing Trend |
| Zn | 30 | µg/l | 60 | 0.0480 | 9.80 | 1.84 | 1.71 | 0.55 | 1.180 | 3.76 | 106 | 0.67 | Decreasing Trend |
| SWQ15 (Bowmanville) | | | | | | | | | | | | | |
| Phosphorus, Total | 30 | µg/l | 69 | 5 | 104 | 21 | 17.200 | 6 | 17 | 35.20 | 390 | 2.01 | Decreasing Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 22 | 0.465 | 1.30 | 0.630 | 0.195 | 0.520 | 0.556 | 0.90 | 33 | 0.90 | No Trend |
| Nitrate as N | 10 | mg/l | 47 | 0.440 | 1.78 | 0.780 | 0.370 | 0.480 | 0.600 | 1.43 | 325 | 2.97 | Decreasing Trend |
| Cl | 250 | mg/l | 69 | 8 | 20 | 12.200 | 3.340 | 8.700 | 11.200 | 17.30 | 473 | 2.44 | Increasing Trend |
| Cu | 5 | µg/l | 49 | 0.053 | 1.80 | 0.395 | 0.287 | 0.200 | 0.300 | 0.65 | 182 | 1.56 | No Trend |
| Cd | 0.1 - 0.5 | µg/l | 45 | 0.009 | 1.13 | 0.155 | 0.175 | 0.100 | 0.100 | 0.28 | 203 | 1.98 | Decreasing Trend |
| Co | 0.9 | µg/l | 46 | 0.027 | 0.89 | 0.233 | 0.168 | 0.100 | 0.200 | 0.30 | 41 | 0.38 | No Trend |
| Fe | 0.3 | mg/l | 56 | 0.063 | 0.32 | 0.115 | 0.046 | 0.074 | 0.105 | 0.17 | 230 | 1.62 | No Trend |
| Ni | 25 | µg/l | 43 | 0.093 | 1.04 | 0.230 | 0.230 | 0.100 | 0.100 | 0.67 | 51 | 0.52 | No Trend |
| Pb | 5 - 25 | µg/l | 42 | 0.100 | 13.30 | 1.250 | 2.320 | 0.100 | 0.700 | 2.92 | 443 | 4.80 | Decreasing Trend |
| Zn | 30 | µg/l | 46 | 0.025 | 2.50 | 0.546 | 0.477 | 0.200 | 0.400 | 1.10 | 270 | 2.55 | Increasing Trend |

| SWQ16 (Bowmanville) | | | | | | | | | | | | | |
|--------------------------------------|-----------|------|----|--------|-------|--------|--------|-------|--------|-------|------|-------|------------------|
| Phosphorus, Total | 30 | µg/l | 44 | 6 | 128 | 21.100 | 25.100 | 6 | 11.500 | 37 | -13 | -0.12 | No Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 0 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | 0 | 0 | No Trend |
| Nitrate as N | 10 | mg/l | 44 | 0.0300 | 1.62 | 0.770 | 0.330 | 0.526 | 0.655 | 1.30 | -148 | -1.49 | No Trend |
| Cl | 250 | mg/l | 44 | 11 | 22.80 | 15.400 | 3.500 | 12 | 14.200 | 20.80 | 216 | 2.17 | Increasing Trend |
| Cu | 5 | µg/l | 44 | 0.100 | 1.50 | 0.434 | 0.295 | 0.20 | 0.35 | 0.80 | 76 | 0.76 | No Trend |
| Cd | 0.1 - 0.5 | µg/l | 44 | 0.100 | 0.20 | 0.105 | 0.021 | 0.10 | 0.10 | 0.10 | -46 | -0.46 | No Trend |
| Co | 0.9 | µg/l | 44 | 0.100 | 0.60 | 0.198 | 0.079 | 0.10 | 0.20 | 0.20 | -90 | -0.9 | Decreasing Trend |
| Fe | 0.3 | mg/l | 44 | 0.037 | 0.23 | 0.083 | 0.044 | 0.05 | 0.07 | 0.13 | -202 | -2.03 | Decreasing Trend |
| Ni | 25 | µg/l | 44 | 0.100 | 0.80 | 0.177 | 0.179 | 0.10 | 0.10 | 0.44 | 310 | 3.13 | Increasing Trend |
| Pb | 5 - 25 | µg/l | 44 | 0.100 | 0.70 | 0.510 | 0.273 | 0.10 | 0.70 | 0.70 | -396 | -4 | Decreasing Trend |
| Zn | 30 | µg/l | 44 | 0.100 | 19.20 | 1.330 | 2.870 | 0.30 | 0.65 | 2.50 | -91 | -0.91 | No Trend |
| SWQ17 (Bowmanville) | | | | | | | | | | | | | |
| Phosphorus, Total | 30 | µg/l | 35 | 5 | 39 | 19.100 | 8.200 | 10.40 | 18 | 29.60 | 116 | 1.63 | No Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 22 | 0.122 | 0.19 | 0.166 | 0.017 | 0.14 | 0.17 | 0.18 | 108 | 3.02 | Decreasing Trend |
| Nitrate as N | 10 | mg/l | 13 | 0.132 | 0.20 | 0.162 | 0.022 | 0.14 | 0.16 | 0.19 | 9 | 0.49 | No Trend |
| Cl | 250 | mg/l | 35 | 0.900 | 1.30 | 1.070 | 0.107 | 0.90 | 1.10 | 1.20 | 15 | 0.20 | No Trend |
| Cu | 5 | µg/l | 12 | 0.113 | 0.56 | 0.303 | 0.161 | 0.12 | 0.27 | 0.53 | 30 | 1.99 | Increasing Trend |
| Cd | 0.1 - 0.5 | µg/l | 9 | 0.008 | 0.76 | 0.328 | 0.260 | 0.11 | 0.26 | 0.70 | 4 | 0.31 | No Trend |
| Co | 0.9 | µg/l | 11 | 0.012 | 1.60 | 0.630 | 0.476 | 0.04 | 0.62 | 1.04 | 11 | 0.78 | No Trend |
| Fe | 0.3 | mg/l | 22 | 0.042 | 0.17 | 0.074 | 0.030 | 0.05 | 0.07 | 0.10 | 83 | 2.31 | Increasing Trend |
| Ni | 25 | µg/l | 8 | 0.285 | 1.18 | 0.600 | 0.360 | 0.30 | 0.47 | 1.16 | 4 | 0.37 | No Trend |
| Pb | 5 - 25 | µg/l | 9 | 0.364 | 4.70 | 1.920 | 1.400 | 0.77 | 1.61 | 3.86 | 8 | 0.73 | No Trend |
| Zn | 30 | µg/l | 14 | 0.315 | 4.76 | 0.87 | 1.14 | 0.363 | 0.52 | 0.96 | 1 | 0 | No Trend |
| SWQ5 (Soper) | | | | | | | | | | | | | |
| Phosphorus, Total | 30 | µg/l | 48 | 6 | 102 | 30.60 | 25 | 6 | 24.50 | 71 | -165 | -1.46 | No Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 3 | 0.990 | 2.75 | 1.80 | 0.89 | 1.13 | 1.68 | 2.54 | -1 | 0 | No Trend |
| Nitrate as N | 10 | mg/l | 45 | 0.810 | 3.95 | 1.66 | 0.79 | 1.06 | 1.36 | 2.87 | 7 | 0.06 | No Trend |
| Cl | 250 | mg/l | 48 | 19.200 | 246 | 33.50 | 33.50 | 20.7 | 25.80 | 37.40 | 59 | 0.52 | No Trend |
| Cu | 5 | µg/l | 48 | 0.200 | 2.4 | 0.76 | 0.55 | 0.2 | 0.60 | 1.56 | 51 | 0.44 | No Trend |

| | | | | | | | | | | | | | |
|--------------------------------------|-----------|------|----|--------|-------|-------|--------|--------|-------|-------|------|-------|------------------|
| Cd | 0.1 - 0.5 | µg/l | 46 | 0.100 | 0.67 | 0.13 | 0.10 | 0.1 | 0.10 | 0.10 | -81 | -0.76 | No Trend |
| Co | 0.9 | µg/l | 47 | 0.100 | 0.71 | 0.26 | 0.12 | 0.2 | 0.20 | 0.44 | 251 | 2.30 | No Trend |
| Fe | 0.3 | mg/l | 48 | 0.031 | 0.41 | 0.14 | 0.09 | 0.0575 | 0.11 | 0.26 | -257 | -2.28 | Decreasing Trend |
| Ni | 25 | µg/l | 46 | 0.100 | 1.1 | 0.30 | 0.29 | 0.1 | 0.20 | 0.80 | 420 | 3.97 | Increasing Trend |
| Pb | 5 - 25 | µg/l | 46 | 0.100 | 8.3 | 0.80 | 1.26 | 0.1 | 0.70 | 0.70 | -329 | -3.10 | Decreasing Trend |
| Zn | 30 | µg/l | 48 | 0.500 | 9.4 | 2.20 | 1.86 | 0.67 | 1.55 | 4.63 | -131 | -1.16 | No Trend |
| SWQ21 (Soper - East) | | | | | | | | | | | | | |
| Phosphorus, Total | 30 | µg/l | 9 | 20 | 100 | 54.30 | 25.90 | 21.6 | 53 | 83.20 | -12 | -1.15 | No Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 0 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | 0 | 0 | No Trend |
| Nitrate as N | 10 | mg/l | 9 | 3.060 | 5.96 | 4.34 | 0.96 | 3.26 | 4.27 | 5.66 | 20 | 1.98 | Increasing Trend |
| Cl | 250 | mg/l | 9 | 27.600 | 37.10 | 33.90 | 2.830 | 31.40 | 34.40 | 36.50 | 18 | 1.77 | Increasing Trend |
| Cu | 5 | µg/l | 9 | 0.200 | 1.10 | 0.68 | 0.303 | 0.20 | 0.80 | 0.94 | 5 | 0.42 | No Trend |
| Cd | 0.1 - 0.5 | µg/l | 9 | 0.100 | 0.20 | 0.11 | 0.033 | 0.10 | 0.10 | 0.12 | -2 | -0.11 | No Trend |
| Co | 0.9 | µg/l | 9 | 0.200 | 0.60 | 0.27 | 0.132 | 0.20 | 0.20 | 0.36 | 13 | 1.26 | No Trend |
| Fe | 0.3 | mg/l | 9 | 0.090 | 0.28 | 0.18 | 0.057 | 0.10 | 0.19 | 0.22 | -10 | -0.94 | No Trend |
| Ni | 25 | µg/l | 9 | 0.100 | 1.20 | 0.39 | 0.390 | 0.10 | 0.20 | 0.88 | 17 | 1.68 | Increasing Trend |
| Pb | 5 - 25 | µg/l | 9 | 0.100 | 0.70 | 0.51 | 0.285 | 0.10 | 0.70 | 0.70 | -18 | -1.80 | Decreasing Trend |
| Zn | 30 | µg/l | 9 | 0.600 | 1.70 | 1.17 | 0.316 | 0.84 | 1.20 | 1.46 | -1 | 0 | No Trend |
| SWQ18 (Soper-East) | | | | | | | | | | | | | |
| Phosphorus, Total | 30 | µg/l | 44 | 6 | 72 | 16.40 | 13.800 | 6 | 10.50 | 33.30 | -55 | -0.55 | No Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 0 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | 0 | 0 | No Trend |
| Nitrate as N | 10 | mg/l | 44 | 0.810 | 2.20 | 1.23 | 0.247 | 1 | 1.17 | 1.52 | -283 | -2.85 | Decreasing Trend |
| Cl | 250 | mg/l | 44 | 8.600 | 23.90 | 19.20 | 3.300 | 17.60 | 19.70 | 21.80 | 366 | 3.70 | Increasing Trend |
| Cu | 5 | µg/l | 44 | 0.100 | 1 | 0.35 | 0.197 | 0.20 | 0.30 | 0.57 | 126 | 1.26 | No Trend |
| Cd | 0.1 - 0.5 | µg/l | 44 | 0.100 | 0.20 | 0.10 | 0.015 | 0.10 | 0.10 | 0.10 | -25 | -0.24 | No Trend |
| Co | 0.9 | µg/l | 44 | 0.100 | 0.40 | 0.21 | 0.068 | 0.10 | 0.20 | 0.27 | 41 | 0.41 | No Trend |
| Fe | 0.3 | mg/l | 44 | 0.010 | 0.30 | 0.06 | 0.059 | 0.02 | 0.04 | 0.12 | -235 | -2.37 | Decreasing Trend |
| Ni | 25 | µg/l | 44 | 0.100 | 0.9 | 0.198 | 0.203 | 0.10 | 0.10 | 0.50 | 354 | 3.57 | Increasing Trend |
| Pb | 5 - 25 | µg/l | 44 | 0.100 | 0.7 | 0.505 | 0.280 | 0.10 | 0.70 | 0.70 | -426 | -4.30 | Decreasing Trend |
| Zn | 30 | µg/l | 44 | 0.100 | 2.3 | 0.560 | 0.494 | 0.20 | 0.40 | 0.97 | -40 | -0.40 | No Trend |

| SWQ19 (Soper-East) | | | | | | | | | | | | | |
|--------------------------------------|-----------|------|----|-------|-------|--------|--------|------|-------|-------|------|-------|------------------|
| Phosphorus, Total | 30 | µg/l | 44 | 6 | 62 | 16.600 | 14.600 | 6 | 10.50 | 36.30 | -1 | 0 | No Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 0 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | 0 | 0 | No Trend |
| Nitrate as N | 10 | mg/l | 44 | 0.790 | 2.28 | 1.410 | 0.293 | 1.18 | 1.36 | 1.77 | 40 | 0.40 | No Trend |
| Cl | 250 | mg/l | 44 | 8.200 | 62.8 | 12.100 | 8.400 | 8.50 | 9.95 | 16.6 | 101 | 1.01 | No Trend |
| Cu | 5 | µg/l | 44 | 0.100 | 2.6 | 0.370 | 0.387 | 0.20 | 0.20 | 0.50 | 154 | 1.55 | No Trend |
| Cd | 0.1 - 0.5 | µg/l | 44 | 0.100 | 0.1 | 0.100 | 0 | 0.10 | 0.10 | 0.10 | 0 | 0 | No Trend |
| Co | 0.9 | µg/l | 44 | 0.100 | 0.5 | 0.207 | 0.073 | 0.10 | 0.20 | 0.30 | -9 | -0.08 | No Trend |
| Fe | 0.3 | mg/l | 44 | 0.023 | 0.436 | 0.072 | 0.068 | 0.03 | 0.05 | 0.13 | -132 | -1.33 | No Trend |
| Ni | 25 | µg/l | 44 | 0.100 | 17.9 | 0.670 | 2.7 | 0.10 | 0.10 | 0.67 | 334 | 3.37 | Increasing Trend |
| Pb | 5 - 25 | µg/l | 44 | 0.100 | 1.5 | 0.523 | 0.315 | 0.10 | 0.70 | 0.70 | -421 | -4.25 | Decreasing Trend |
| Zn | 30 | µg/l | 44 | 0.100 | 3.5 | 0.820 | 0.870 | 0.20 | 0.45 | 2.10 | 14 | 0.13 | No Trend |
| SWQ20 (Soper-East) | | | | | | | | | | | | | |
| Phosphorus, Total | 30 | µg/l | 6 | 6 | 14 | 7.700 | 3.200 | 6 | 6 | 11 | -5 | -0.77 | No Trend |
| Nitrate (Total, Unfiltered Reactive) | 10 | mg/l | 0 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | 0 | 0 | No Trend |
| Nitrate as N | 10 | mg/l | 6 | 0.750 | 1.08 | 0.93 | 0.14 | 0.78 | 0.945 | 1.07 | -15 | -2.63 | Decreasing Trend |
| Cl | 250 | mg/l | 6 | 6 | 9.80 | 7.54 | 1.69 | 6.10 | 6.900 | 9.60 | 7 | 1.13 | No Trend |
| Cu | 5 | µg/l | 6 | 0.200 | 1.20 | 0.53 | 0.39 | 0.25 | 0.350 | 1 | -4 | -0.57 | No Trend |
| Cd | 0.1 - 0.5 | µg/l | 6 | 0.100 | 0.10 | 0.1 | 0 | 0.10 | 0.100 | 0.10 | 0 | 0 | No Trend |
| Co | 0.9 | µg/l | 6 | 0.100 | 0.20 | 0.15 | 0.06 | 0.10 | 0.150 | 0.20 | -9 | -1.56 | No Trend |
| Fe | 0.3 | mg/l | 6 | 0.024 | 0.13 | 0.07 | 0.04 | 0.03 | 0.075 | 0.11 | -11 | -1.88 | Decreasing Trend |
| Ni | 25 | µg/l | 6 | 0.100 | 0.60 | 0.33 | 0.21 | 0.10 | 0.350 | 0.55 | 6 | 0.96 | No Trend |
| Pb | 5 - 25 | µg/l | 6 | 0.100 | 10.20 | 2.08 | 4 | 0.10 | 0.700 | 5.45 | -5 | -0.78 | No Trend |
| Zn | 30 | µg/l | 6 | 0.200 | 1.60 | 0.67 | 0.59 | 0.20 | 0.400 | 1.40 | -6 | -0.94 | No Trend |

Table B-9: Statistical Results for Select Surface Water Quality Sites within CLOSPA

All analysis was completed with AquaChem Water Quality software.

B1.6 GROUNDWATER QUALITY DATA ANALYSIS AND REPORTING

B1.6.1 Data Compilation

Groundwater quality data may be available from a wide variety of sources, including:

- The Provincial Groundwater Monitoring Network (PGMN);
- Private well sampling;
- Municipal water sampling programs;
- Health departments; and
- Other groundwater studies.

B1.6.2 Data Analysis

The assemblage and integration of information that will provide an understanding of groundwater quality on watershed basis can be performed a number of ways, including:

- The assemblage of GIS layers;
- The construction of binary plots;
- The construction of maps and cross sections;
- The construction of vertical and horizontal iso-chemical contour maps;
- The construction of groundwater quality diagrams (e.g., Durov, Piper, Stiff, Rose);
- The construction of chemical concentration versus time plots;
- The preparation of tables that compare water quality concentrations to water quality criteria (e.g., Ontario Drinking Water Standards, Provincial Water Quality Standards); and
- The use of statistical methods.

Parameters that exceed the standard can be highlighted, as some parameters naturally exceed water quality standards. Naturally elevated parameters can be present due to the geological materials in the area, the recharge environment, or other factors.

B1.6.3 Analysis of Trends at Each Monitoring Well

Time versus concentration plots can help determine whether levels of water quality are changing. Time-concentration plots are generated from water quality data for one parameter, usually in one monitoring well, with time across the x-axis, and the concentration for that parameter along the y-axis. Statistical trend analysis packages (e.g., packages built into Excel) can be used to determine if there is a trend. See Error! Reference source not found..

Alternatively, the data can be visually interpreted to determine whether there is a trend. Trends usually occur over a longer term, though there may be a blip or short-term spike in concentration indicating a short-term event, such as a spill or controlled release into the environment. Trends can also occur seasonally or cyclically. Seasonal or cyclic trends occur where water quality fluctuates through seasons or through wet or dry years.

Where water quality impairments have been identified in a watershed (i.e., concerns, known contamination), the parameters typical for those impairments can also be evaluated through time-concentration plots to determine whether the trends are increasing or decreasing. Trend analysis can provide an indication of contamination, changes in groundwater recharge, a connection to surface water, or general changes within an aquifer. Significant increasing or decreasing trends should be identified in the individual monitoring wells. By doing this, we can identify areas where water quality is influenced by surface activities, including precipitation, and therefore may be more vulnerable to surface activities.

| Station ID | Sampling Date | 1,4-Dichlorobenzene | Al | Cl | Fe | Mn | Na | Pb | Se | Zn | Colour | Carbon; dissolved organic | Hardness as CaCO ₃ | Turbidity |
|--|---------------|---------------------|----------|----------|----------|-----------|---------|---------|---------|-----------|--------|---------------------------|-------------------------------|-----------|
| Ontario Drinking Water Standards (MAC, AO) | | 1 µg/L | 0.1 mg/L | 250 mg/L | 0.3 mg/L | 0.05 mg/L | 20 mg/L | 10 µg/L | 10 µg/L | 5000 µg/L | 5 TCU | 5 mg/L | 80 mg/L | 5 NTU |
| W0000263-1 | 31/07/2003 | 0.05 | 0.0319 | 268 | 0.0490 | 0.0032 | 203 | 0.05 | 0 | 0.2 | | 1.3 | | |
| W0000263-1 | 15/06/2004 | | 0.0079 | 388 | 0.0021 | <0.0001 | 196 | <0.7 | <2 | 1.9 | 2 | | 345 | 0.087 |
| W0000263-1 | 09/07/2004 | | 0.0065 | 232 | <0.0002 | <0.0001 | 171 | <0.7 | <2 | 2.1 | <1 | | 219 | 0.072 |
| W0000263-1 | 27/06/2005 | | <0.0007 | 241 | 0 | <0.0001 | 194 | <0.7 | <2 | 0 | 2 | | 221 | 0.080 |
| W0000263-1 | 18/10/2005 | | <0.0007 | 221 | 0 | <0.0001 | 170 | <0.7 | <2 | 0 | 1 | | 174 | 0.080 |
| W0000263-1 | 15/06/2006 | | <0.0007 | 334 | 0.0056 | <0.0001 | 233 | <0.7 | <2 | 0.4 | 1 | | 231 | 0.135 |
| W0000263-1 | 30/10/2006 | | <0.0007 | 331 | 0.0060 | <0.0001 | 221 | <0.7 | <2 | 2 | 3 | | 259 | 0.105 |
| W0000263-1 | 11/05/2007 | | | 145 | 0.0436 | 0.0022 | 132 | 0.1 | 0.3 | 0.8 | 2 | | 191 | 0.366 |
| W0000263-1 | 29/05/2007 | | | 157 | 0.0070 | <0.0001 | 134 | <0.7 | <2 | 0.3 | 11 | | 183 | 0.086 |
| W0000263-1 | 29/05/2008 | | 0.0177 | 122 | 0.0229 | 0.0011 | 122 | <0.7 | 0.1 | 0.8 | 3 | | 155 | 0.250 |
| W0000263-1 | 10/06/2008 | | 0.0123 | 224 | 0.0040 | 0.00097 | 145 | 0.04 | 0 | 0.9 | | 1.3 | 192 | |
| W0000040-1 | 14/08/2003 | 0.05 | 0.0011 | 238 | 0.0010 | 0.00022 | 106 | 0.04 | 1 | 2.2 | | 2.2 | 628 | |
| W0000040-1 | 23/06/2004 | | 0.0021 | 168 | 0.0036 | <0.0001 | 70.4 | <0.7 | 3 | 2.4 | 2 | | 475 | 0.069 |
| W0000040-1 | 20/10/2004 | | 0.0034 | 462 | 0.0023 | <0.0001 | 155 | <0.7 | <2 | 1.8 | <1 | | 734 | 0.119 |
| W0000040-1 | 11/03/2005 | | <0.0007 | 555 | 0.0100 | <0.0001 | 229 | <0.7 | <2 | 0 | 3 | | 751 | 0.080 |
| W0000040-1 | 07/06/2005 | | <0.0007 | 318 | 0.0100 | <0.0001 | 113 | <0.7 | <2 | 0 | 1 | | 603 | 0.080 |
| W0000040-1 | 11/01/2006 | | <0.0007 | 115 | 0.0091 | <0.0001 | 82.8 | <0.7 | <2 | 0.2 | <1 | | 403 | 0.083 |
| W0000040-1 | 06/12/2006 | | <0.0007 | 119 | 0.0061 | <0.0001 | 79.7 | <0.7 | <2 | <0.2 | 1 | | 382 | 0.103 |
| W0000040-1 | 13/06/2007 | | | 133 | 0.0149 | <0.0001 | 70.4 | <0.7 | <2 | 0.4 | 1 | | 381 | 0.068 |
| W0000040-1 | 20/11/2007 | | | 667 | 0.0077 | 0.0020 | 271 | <0.7 | 0.7 | 0.8 | 3 | | 779 | 0.245 |
| W0000040-1 | 06/02/2008 | | 0.0009 | 111 | 0.0002 | 0.0001 | 84.4 | <0.7 | 0.3 | 0.6 | 2 | | 336 | 0.121 |
| W0000040-1 | 18/09/2008 | | 0.0008 | 110 | 0.0060 | 0.0026 | 91.3 | 0.11 | 1 | 1.1 | | 1.3 | 213 | |
| W0000044-2 | 18/09/2002 | 0.05 | 0.0013 | 19.3 | 0.1840 | 0.1330 | 7.8 | 0.24 | 0 | 613 | | 1.4 | 306 | |

| Station ID | Sampling Date | 1,4-Dichlorobenzene | Al | Cl | Fe | Mn | Na | Pb | Se | Zn | Colour | Carbon; dissolved organic | Hardness as CaCO3 | Turbidity |
|--|---------------|---------------------|----------|----------|----------|-----------|---------|---------|---------|-----------|--------|---------------------------|-------------------|-----------|
| Ontario Drinking Water Standards (MAC, AO) | | 1 µg/L | 0.1 mg/L | 250 mg/L | 0.3 mg/L | 0.05 mg/L | 20 mg/L | 10 µg/L | 10 µg/L | 5000 µg/L | 5 TCU | 5 mg/L | 80 mg/L | 5 NTU |
| W0000044-2 | 07/06/2004 | | 0.0075 | 19.3 | 0.2220 | 0.1030 | 7.6 | <0.7 | <2 | 776 | <1 | | 304 | 0.901 |
| W0000044-2 | 10/06/2004 | | 0.0037 | 19 | 0.5050 | 0.1190 | 7.6 | <0.7 | <2 | 799 | 5 | | 300 | 2.150 |
| W0000044-2 | 16/06/2005 | | <0.0007 | 19.8 | 0.5500 | 0.1000 | 7.4 | <0.7 | <2 | 880 | 36 | | 297 | 3.460 |
| W0000044-2 | 26/10/2005 | | 0.0300 | 19.7 | 0.5500 | 0.1500 | 7.7 | <0.7 | <2 | 960 | 18 | | 309 | 3.730 |
| W0000044-2 | 06/08/2006 | | 0.165 | 18 | 0.2300 | 0.0607 | 7.8 | 11.9 | <2 | 2100 | 28 | | 254 | 25.400 |
| W0000044-2 | 19/10/2006 | | 9.4400 | 19.8 | 18 | 0.5760 | 7.8 | 193 | 2 | 51300 | 7 | | 273 | 1.540 |
| W0000044-2 | 10/09/2008 | | 0.0213 | 21.7 | 0.2920 | 0.1340 | 7.5 | 4.17 | 0 | 1667.4 | | 1.3 | 305 | |
| W0000264-2 | 08/12/2003 | 2 | 0.2380 | 34 | 1.0200 | 0.0400 | 129 | 9.69 | 0 | 20.5 | | 0.2 | 393 | |
| W0000264-2 | 10/05/2004 | | <0.0007 | 2.4 | <0.0002 | 0.0038 | 10.5 | <0.7 | <2 | 13.6 | 3 | | 149 | 0.176 |
| W0000264-2 | 16/06/2004 | | 0.1480 | 4.44 | 0.0471 | 0.0037 | 30 | <0.7 | <2 | 3.1 | 16 | | 97.5 | 4.42 |
| W0000264-2 | 20/07/2004 | | 0.0977 | 3.42 | 0.0537 | 0.0047 | 21 | <0.7 | <2 | 19.9 | 6 | | 118 | 0.469 |
| W0000264-2 | 18/10/2006 | | 0.6050 | 5.33 | 0.247 | 0.0032 | 3.36 | <0.7 | <2 | 1.6 | 28 | | 162 | 7.48 |
| W0000264-2 | 30/05/2007 | | | 6.7 | 0.0443 | 0.0017 | 3.5 | <0.7 | <2 | 0.7 | 6 | | 183 | 0.399 |
| W0000264-2 | 16/09/2008 | | 0.2410 | 5.8 | 0.196 | 0.022 | 3.16 | 0.4 | 0 | 3 | | 0.2 | 171 | |
| W0000167-1 | 28/11/2002 | 0.05 | 0.0019 | 20.9 | 1.310 | 0.189 | 12.6 | 0 | 0 | 1 | | 6 | 329 | |
| W0000167-1 | 21/06/2004 | | <0.0007 | 22.3 | 1.090 | 0.167 | 14.4 | <0.7 | <2 | 0.3 | 19 | | 291 | 8.02 |
| W0000167-1 | 09/08/2004 | | <0.0007 | 18.1 | 0.991 | 0.186 | 16.1 | <0.7 | <2 | 1.9 | 24 | | 325 | 4.16 |
| W0000167-1 | 26/05/2005 | | <0.0007 | 24.1 | 1.150 | 0.240 | 11 | <0.7 | <2 | 0 | 19 | | 283 | 4.63 |
| W0000167-1 | 19/10/2005 | | 0 | 16.1 | 1.680 | 0.310 | 12.6 | <0.7 | <2 | 0 | 20 | | 312 | 7.34 |
| W0000167-1 | 06/06/2006 | | 0.0581 | 22.6 | 4.520 | 0.283 | 12.9 | <0.7 | <2 | 0.7 | 51 | | 284 | 48.8 |
| W0000167-1 | 16/10/2006 | | <0.0007 | 17.6 | 2.490 | 0.278 | 13.3 | <0.7 | <2 | 0.2 | 23 | | 298 | 8.13 |
| W0000167-1 | 11/07/2007 | | | 11.8 | 3.670 | 0.283 | 10.9 | <0.7 | <2 | 0.2 | 30 | | 268 | 19.2 |
| W0000167-1 | 06/12/2007 | | | 15.7 | 1.740 | 0.234 | 11.5 | <0.7 | <2 | 0.5 | 37 | | 279 | 13.1 |
| W0000167-1 | 27/05/2008 | | 0.0014 | 32.7 | 1.170 | 0.192 | 10.9 | <0.7 | 0.3 | 0.1 | 57 | | 296 | 2.65 |
| W0000167-1 | 15/09/2008 | | 0.00158 | 26.1 | 1.360 | 0.218 | 12.9 | 0 | 0 | 4.9 | | 4.9 | 310 | |

| Station ID | Sampling Date | 1,4-Dichlorobenzene | Al | Cl | Fe | Mn | Na | Pb | Se | Zn | Colour | Carbon; dissolved organic | Hardness as CaCO ₃ | Turbidity |
|--|---------------|---------------------|----------|----------|----------|-----------|---------|---------|---------|-----------|--------|---------------------------|-------------------------------|-----------|
| Ontario Drinking Water Standards (MAC, AO) | | 1 µg/L | 0.1 mg/L | 250 mg/L | 0.3 mg/L | 0.05 mg/L | 20 mg/L | 10 µg/L | 10 µg/L | 5000 µg/L | 5 TCU | 5 mg/L | 80 mg/L | 5 NTU |
| W0000049-1 | 24/10/2002 | 0.05 | 0.0003 | 2.4 | 0.311 | 0.008 | 276 | 0 | 0 | 0.2 | | 0.7 | 2100 | |
| W0000049-1 | 29/06/2004 | | 0.004 | 2.6 | 0.465 | 0.007 | 11.9 | <0.7 | <2 | 0.2 | 3 | | 153 | 1.96 |
| W0000049-1 | 14/09/2004 | | 0.0012 | 2.4 | 0.476 | 0.007 | 11.9 | <0.7 | <2 | 1.7 | 12 | | 152 | 1.18 |
| W0000049-1 | 28/06/2005 | | <0.0007 | 2.9 | 0.630 | 0.010 | 12.3 | <0.7 | <2 | 0 | 18 | | 160 | 1.09 |
| W0000049-1 | 11/07/2005 | | <0.0007 | 2.8 | 0.480 | 0.010 | 12.6 | <0.7 | <2 | 0 | 7 | | 160 | 0.81 |
| W0000049-1 | 11/01/2006 | | <0.0007 | 2.9 | 0.409 | 0.007 | 12.6 | <0.7 | <2 | <0.2 | 4 | | 159 | 2.29 |
| W0000049-1 | 15/06/2006 | | <0.0007 | 2.57 | 0.551 | 0.0084 | 12.8 | <0.7 | <2 | <0.2 | 8 | | 158 | 2.06 |
| W0000049-1 | 29/05/2007 | | | 2.9 | 0.618 | 0.0116 | 11.9 | <0.7 | <2 | <0.2 | 11 | | 161 | 1.55 |
| W0000049-1 | 15/11/2007 | | | 3.2 | 0.500 | 0.0096 | 11.4 | <0.7 | <2 | <0.2 | 7 | | 163 | 1.05 |
| W0000049-1 | 29/05/2008 | | 0.0003 | 3.5 | 0.468 | 0.0104 | 11 | <0.7 | <2 | 0.2 | 8 | | 169 | 1.53 |
| W0000049-1 | 18/09/2008 | | 0.0001 | 4.1 | 0.408 | 0.0105 | 10.9 | 0.02 | 0 | 0.2 | | 0.5 | 173 | |
| W0000262-1 | 29/07/2003 | 0.05 | 0.0024 | 9.3 | 0.021 | 0.0498 | 6.8 | 0.03 | 0 | 1 | | 0.4 | 300 | |
| W0000262-1 | 15/09/2008 | | 0.0016 | 26.1 | 1.360 | 0.2180 | 12.9 | 0.02 | 0 | 4.9 | | 4.9 | 310 | |
| W0000261-1 | 31/07/2003 | 0.05 | 0.0012 | 1.3 | 0 | 0.0022 | 3 | 0.02 | 1 | 0.1 | | 0.7 | | |
| W0000261-1 | 28/05/2007 | | | 1.9 | 0.011 | <0.0001 | 2.5 | <0.7 | <2 | 0.3 | 8 | | 227 | 0.09 |
| W0000261-1 | 15/09/2008 | | 0.0012 | 1.3 | 0.006 | 0.0012 | 2.5 | 0.01 | 0 | 0.5 | | 0.4 | 229 | |
| W0000166-1 | 09/11/2003 | 0.05 | 0.0004 | 2.3 | 0.454 | 0.0187 | 6.8 | 0 | 0 | 1.1 | | 0.6 | 181 | |
| W0000166-1 | 21/06/2004 | | 0.0008 | 2.4 | 0.396 | 0.0155 | 7.8 | <0.7 | <2 | <0.2 | 6 | | 169 | 2.28 |
| W0000166-1 | 09/08/2004 | | 0.0028 | 2.5 | 0.342 | 0.0154 | 7.9 | <0.7 | <2 | 1.8 | 2 | | 167 | 1.43 |
| W0000166-1 | 06/09/2005 | | <0.0007 | 2.7 | 0.440 | 0.0200 | 8.2 | <0.7 | <2 | 0 | 1 | | 165 | 2.73 |
| W0000166-1 | 19/10/2005 | | <0.0007 | 2.4 | 0.420 | 0.0200 | 8.6 | <0.7 | <2 | 0 | 3 | | 167 | 1.81 |
| W0000166-1 | 06/06/2006 | | <0.0007 | 2.1 | 0.318 | 0.0152 | 8.5 | <0.7 | <2 | <0.2 | 2 | | 157 | 1.64 |
| W0000166-1 | 11/09/2006 | | <0.0007 | 2.7 | 0.429 | 0.0202 | 9 | <0.7 | <2 | <0.2 | 10 | | 162 | 1.77 |
| W0000166-1 | 19/06/2007 | | | 2.9 | 0.415 | 0.0182 | 8.6 | <0.7 | <2 | <0.2 | 3 | | 159 | 1.21 |
| W0000166-1 | 19/11/2007 | | | 2.8 | 0.445 | 0.0209 | 8.8 | <0.7 | <2 | <0.2 | 7 | | 164 | 1.81 |

| Station ID | Sampling Date | 1,4-Dichlorobenzene | Al | Cl | Fe | Mn | Na | Pb | Se | Zn | Colour | Carbon; dissolved organic | Hardness as CaCO3 | Turbidity |
|--|---------------|---------------------|----------|----------|----------|-----------|---------|---------|---------|-----------|--------|---------------------------|-------------------|-----------|
| Ontario Drinking Water Standards (MAC, AO) | | 1 µg/L | 0.1 mg/L | 250 mg/L | 0.3 mg/L | 0.05 mg/L | 20 mg/L | 10 µg/L | 10 µg/L | 5000 µg/L | 5 TCU | 5 mg/L | 80 mg/L | 5 NTU |
| W0000166-1 | 06/03/2008 | | 0.0004 | 2.5 | 0.422 | 0.0204 | 8.4 | <0.7 | 0.1 | 0.6 | 3 | | 161 | 0.48 |
| W0000166-1 | 17/09/2008 | | 0.0001 | 2.7 | 0.357 | 0.0167 | 8.76 | 0.01 | 0 | 0.4 | | 0.5 | 153 | |
| W0000264-3 | 08/12/2003 | 0.05 | 0.0012 | 3 | 0.137 | 0.0155 | 26.4 | 0.03 | 0 | 3.3 | | 0.8 | 165 | |
| W0000264-3 | 16/09/2008 | | 0.0008 | 2 | 0.174 | 0.0094 | 2.94 | 0.01 | 0 | 0.6 | | 0.1 | 190 | |
| W0000042-1 | 28/11/2002 | 0.05 | 0.0006 | 84.4 | 0.015 | 0.0040 | 33.2 | 0.07 | 0 | 0.7 | | 0.5 | 248 | |
| W0000042-1 | 23/06/2004 | | <0.0007 | 93.2 | 0.018 | 0.0016 | 36.5 | <0.7 | <2 | 2.9 | <1 | | 266 | 0.157 |
| W0000042-1 | 20/10/2004 | | <0.0007 | 82.1 | 0.016 | 0.0041 | 33.4 | <0.7 | <2 | 1.8 | <1 | | 262 | 0.094 |
| W0000042-1 | 11/07/2005 | | <0.0007 | 67.5 | 0.010 | 0 | 26.9 | <0.7 | <2 | 0 | <1 | | 262 | 0.080 |
| W0000042-1 | 06/09/2005 | | <0.0007 | 70.9 | 0.020 | 0 | 29 | <0.7 | <2 | 0 | 1 | | 255 | 0.130 |
| W0000042-1 | 11/09/2006 | | <0.0007 | 74.4 | 0.015 | 0.0012 | 30.9 | <0.7 | <2 | 0.4 | <1 | | 266 | 0.100 |
| W0000042-1 | 06/12/2006 | | <0.0007 | 54.9 | 0.014 | 0.0026 | 23.8 | <0.7 | <2 | <0.2 | <1 | | 250 | 0.218 |
| W0000042-1 | 13/06/2007 | | | 76.8 | 0.027 | 0.0019 | 31.2 | <0.7 | <2 | 0.3 | 1 | | 261 | 0.154 |
| W0000042-1 | 14/11/2007 | | | 76.2 | 0.034 | 0.0041 | 33.5 | <0.7 | <2 | <0.2 | <1 | | 261 | 0.214 |
| W0000042-1 | 06/02/2008 | | 0.0009 | 63.5 | 0.012 | 0.0023 | 28.3 | <0.7 | 0.2 | 0.5 | 1 | | 252 | 0.193 |
| W0000042-1 | 17/09/2008 | | <0.0010 | 75 | 0.023 | 0.0068 | 30.1 | 0.14 | 2 | 0.6 | | 0.5 | 255 | |
| W0000168-1 | 21/11/2002 | 0.05 | 0.0018 | 13.1 | 0.124 | 0.0135 | 17.6 | 0.11 | 0 | 0.5 | | 1.1 | 177 | |
| W0000168-1 | 07/12/2004 | | 0.0056 | 11.8 | <0.0002 | 0.0110 | 17.4 | <0.7 | <2 | 1 | 5 | | 135 | 0.164 |
| W0000168-1 | 10/12/2004 | | 1.8900 | 11.3 | 1.170 | 0.0170 | 17.7 | <0.7 | <2 | 5.3 | 2 | | 141 | 96 |
| W0000044-3 | 11/05/2002 | 0.05 | 0.0006 | 360 | 0.084 | 0.0204 | 192 | 1.15 | 2 | 487 | | 3 | 156 | |
| W0000044-3 | 07/06/2004 | | 0.0105 | 83.3 | 0.001 | <0.0001 | 60.2 | <0.7 | <2 | 98.4 | 3 | | 21.4 | 0.228 |
| W0000044-3 | 10/06/2004 | | 0.4290 | 71.8 | 0.575 | 0.0227 | 60.2 | 1.8 | <2 | 275 | 3 | | 28.5 | 1.380 |
| W0000044-3 | 16/06/2005 | | <0.0007 | 58.1 | 0.010 | 0 | 57.1 | <0.7 | <2 | 40 | 4 | | 23.3 | 0.620 |
| W0000044-3 | 06/08/2006 | | 0.613 | 51.2 | 0.375 | 0.0115 | 60.5 | <0.7 | <2 | 66.7 | 124 | | 49.6 | 265 |
| W0000044-3 | 19/10/2006 | | 7.51 | 53.5 | 6.55 | 0.077 | 56.4 | 13.1 | <2 | 1250 | 30 | | 35.5 | 34 |
| W0000044-3 | 10/09/2008 | | 0.0149 | 52.4 | 0.036 | 0.0017 | 55 | 2.42 | 0 | 53.4 | | 1.5 | 14.2 | |

| Station ID | Sampling Date | 1,4-Dichlorobenzene | Al | Cl | Fe | Mn | Na | Pb | Se | Zn | Colour | Carbon; dissolved organic | Hardness as CaCO3 | Turbidity |
|--|---------------|---------------------|----------|----------|----------|-----------|---------|---------|---------|-----------|--------|---------------------------|-------------------|-----------|
| Ontario Drinking Water Standards (MAC, AO) | | 1 µg/L | 0.1 mg/L | 250 mg/L | 0.3 mg/L | 0.05 mg/L | 20 mg/L | 10 µg/L | 10 µg/L | 5000 µg/L | 5 TCU | 5 mg/L | 80 mg/L | 5 NTU |
| W0000043-3 | 11/05/2002 | 0.05 | 0.0090 | 100 | 0.010 | 0.0150 | 69.4 | 0.53 | 1 | 133 | | 85 | 91 | |
| W0000043-3 | 07/06/2004 | | 0.0073 | 368 | 0.046 | 0.0141 | 195 | <0.7 | <2 | 225 | 1 | | 147 | 0.177 |
| W0000043-3 | 10/06/2004 | | 0.0206 | 358 | 0.109 | 0.0160 | 198 | 2.7 | <2 | 485 | 3 | | 153 | 0.554 |
| W0000043-3 | 11/03/2005 | | <0.0007 | 366 | 0.080 | 0.0100 | 198 | <0.7 | <2 | 350 | 5 | | 152 | 0.220 |
| W0000043-3 | 16/06/2005 | | 0.14 | 361 | 0.740 | 0.1100 | 199 | <0.7 | <2 | 20 | 3 | | 151 | 0.250 |
| W0000043-3 | 06/08/2006 | | <0.0007 | 363 | 0.066 | 0.0067 | 212 | <0.7 | <2 | 123 | 2 | | 151 | 0.580 |
| W0000043-3 | 30/10/2006 | | 0.0376 | 359 | 0.102 | 0.0097 | 209 | <0.7 | <2 | 397 | 3 | | 156 | 3.210 |
| W0000043-3 | 11/05/2007 | | | 341 | 0.094 | 0.0152 | 196 | 1.9 | <2 | 270 | 2 | | 141 | 0.177 |
| W0000043-3 | 19/06/2007 | | | 374 | 0.094 | 0.0137 | 209 | 0.8 | <2 | 460 | 3 | | 159 | 0.170 |
| W0000043-3 | 26/05/2008 | | 0.0406 | 342 | 0.145 | 0.0132 | 196 | 1 | 11.4 | 420 | 1 | | 140 | 0.575 |
| W0000043-3 | 17/09/2008 | | 1.19 | 348 | 1.120 | 0.0465 | 194 | 6.65 | 5 | 1750 | | 1.1 | 165 | |
| W0000041-1 | 21/11/2002 | 0.05 | 0.0009 | 40.8 | 0.199 | 0.0195 | 38.4 | 0.04 | 2 | 0.6 | | 0.7 | 83.5 | |
| W0000041-1 | 13/09/2004 | | 0.0012 | 31.7 | 0.406 | 0.0325 | 29.5 | <0.7 | <2 | 1.7 | 9 | | 99.3 | 0.578 |
| W0000041-1 | 07/12/2004 | | 0.0016 | 33.5 | 0.289 | 0.0258 | 30.4 | <0.7 | <2 | <0.2 | 6 | | 96.2 | 0.441 |
| W0000041-1 | 06/06/2005 | | <0.0007 | 34.1 | 0.420 | 0.0500 | 29 | <0.7 | <2 | 0 | 7 | | 104 | 0.890 |
| W0000041-1 | 20/10/2005 | | <0.0007 | 33.1 | 0.410 | 0.0400 | 29.2 | <0.7 | <2 | <0.2 | 7 | | 104 | 0.580 |
| W0000041-1 | 16/10/2006 | | <0.0007 | 45.3 | 0.326 | 0.0251 | 33.2 | <0.7 | <2 | <0.2 | 3 | | 92.5 | 0.394 |
| W0000041-1 | 06/12/2006 | | <0.0007 | 56.4 | 0.202 | 0.0169 | 37.6 | <0.7 | <2 | <0.2 | 5 | | 91.2 | 0.405 |
| W0000041-1 | 11/07/2007 | | | 33.8 | 0.601 | 0.0432 | 29.5 | <0.7 | <2 | <0.2 | 5 | | 106 | 0.823 |
| W0000041-1 | 06/12/2007 | | | 38.7 | 0.217 | 0.0243 | 30.6 | <0.7 | <2 | <0.2 | 3 | | 92.9 | 0.286 |
| W0000041-1 | 26/05/2008 | | <0.0002 | 37.3 | 0.154 | 0.0287 | 30.5 | <0.7 | 0.6 | 1.5 | 1 | | 94.3 | 0.286 |
| W0000041-1 | 10/06/2008 | | 0.00004 | 36.9 | 0.860 | 0.0608 | 30.5 | 0.01 | 0 | 0.4 | | 1.1 | 104 | |
| W0000265-2 | 08/12/2003 | 0.05 | 0.0028 | 3.2 | 0.102 | 0.0130 | 28.4 | 0.02 | 0 | 1.8 | | 1.8 | 94.5 | |
| W0000265-2 | 10/05/2004 | | <0.0007 | 3.1 | 0.148 | 0.0161 | 24.9 | <0.7 | <2 | 2.4 | 6 | | 110 | 0.160 |
| W0000265-2 | 16/06/2004 | | 0.0047 | 3.2 | 0.131 | 0.0126 | 25.3 | <0.7 | <2 | 0.2 | 7 | | 106 | 0.129 |

| Station ID | Sampling Date | 1,4-Dichlorobenzene | Al | Cl | Fe | Mn | Na | Pb | Se | Zn | Colour | Carbon; dissolved organic | Hardness as CaCO ₃ | Turbidity |
|--|---------------|---------------------|----------|----------|----------|-----------|---------|---------|---------|-----------|--------|---------------------------|-------------------------------|-----------|
| Ontario Drinking Water Standards (MAC, AO) | | 1 µg/L | 0.1 mg/L | 250 mg/L | 0.3 mg/L | 0.05 mg/L | 20 mg/L | 10 µg/L | 10 µg/L | 5000 µg/L | 5 TCU | 5 mg/L | 80 mg/L | 5 NTU |
| W0000265-2 | 19/07/2004 | | <0.0007 | 3.1 | 0.113 | 0.0120 | 24.5 | <0.7 | <2 | 2 | 8 | | 108 | 0.136 |
| W0000265-2 | 15/06/2005 | | <0.0007 | 3.2 | 0.140 | 0.0100 | 24.4 | 0 | <2 | 0 | 6 | | 110 | 0.160 |
| W0000265-2 | 24/10/2005 | | <0.0007 | 3.0 | 0.160 | 0.0200 | 24.4 | <0.7 | <2 | 0 | 6 | | 110 | 0.170 |
| W0000265-2 | 06/07/2006 | | <0.0007 | 2.8 | 0.146 | 0.0150 | 24.7 | <0.7 | <2 | 0.5 | 6 | | 108 | 0.214 |
| W0000265-2 | 18/10/2006 | | <0.0007 | 3.2 | 0.140 | 0.0151 | 25 | <0.7 | <2 | <0.2 | 6 | | 110 | 0.179 |
| W0000265-2 | 30/05/2007 | | | 3.6 | 0.155 | 0.0161 | 24.7 | <0.7 | <2 | <0.2 | 6 | | 110 | 0.149 |
| W0000265-2 | 13/11/2007 | | | 3.0 | 0.151 | 0.0167 | 24.6 | <0.7 | <2 | 0.2 | 5 | | 110 | 0.197 |
| W0000265-2 | 28/05/2008 | | 0.0011 | 3 | 0.130 | 0.0149 | 24.5 | <0.7 | <2 | 0.1 | 5 | | 108 | 0.237 |
| W0000265-2 | 16/09/2008 | | 0.00107 | 3.1 | 0.098 | 0.0128 | 24.3 | 0.01 | 0 | 0.3 | | 0.9 | 104 | |

Table B-10: List of all Groundwater Quality Samples within CLOSPA with Parameters that have Exceeded the ODWS at Least once

All instances are highlighted in red.

B1.6.4 Aquifer Characterization

Groundwater quality data was also analyzed on a watershed basis to look for larger-scale trends in water quality. Monitoring wells from similar aquifer units can be grouped to determine the typical maximum, minimum, and average water quality ranges for the aquifer units. Where little information is available to determine whether monitoring wells are in the same aquifer, water quality data can be compared through Piper diagrams, Stiff diagrams, Rose diagrams, and other geochemistry tools to determine whether water samples are of a similar nature, and potentially of similar origin.

B1.7 LIMITATIONS: DATA, ASSUMPTIONS, AND METHODS

B1.7.1 Data

Database management that relates to the structure or approach was developed for each of the conservation authority partners and the CTC Region to manage data. Currently, a three-database system is being considered within the overall database management system. This system includes:

- Internal relational databases that house aquatic ecosystem and stream survey information conducted by CLOSPA;
- The Oak Ridges Moraine Groundwater Program database that includes subsurface information (e.g., boreholes, wells, water levels, chemistry); and
- The contaminant inventory database, to be provided by the Province.

Data that are undergoing refinement have been identified for source protection planning purposes and are summarized in **Table B-11**.

| Identified Data that is undergoing Refinement (not available at the time of reporting) | | | |
|--|--|-----------------|---|
| Watershed Characterization | | | |
| Component | Data Set Name or Source | Data Problem | Comment |
| GIS Database | CLOCA/external data sources | Requires update | Internal GIS data, grids, shape file reorganization. Metadata tracking system to be developed. |
| Rating Tables within Hydrologic Database | CLOCA — Engineering department hydraulic data | Requires update | Updated for WSC sites annually. CLOCA sites to be surveyed and calibrated. Need to be generated for 2005 - 2006. |
| Integrated Hydrologic Database | CLOCA's hydrologic data | Requires update | Data currently exists in various formats. Need to develop a consistent format and relational database to maintain data relating to climate, rating curves, water levels, streamflow, spot baseflow, and water quality measurements. |
| Oak Ridges Moraine Groundwater Program Hydrogeologic Database | Various data sources | Requires update | Not all monitoring locations or data entered—continually being updated with various data sets. Database management is required. Multi-user access to be applied over a networked environment. |
| Water Quality/Benthic Database | Provincial OBBN Historical studies and reports | Requires update | Data, in general, has not been QC'd. |

| Knowledge Gaps |
|--|
| <ul style="list-style-type: none">• Continued groundwater level and chemistry monitoring and analysis involving both PGMN wells and municipal partner monitoring wells (where data is provided).• Low-flow streamflow surveys (quality and quantity) to characterize discharge zones and associated water quality. These surveys are also useful to delineate zones that may be impacted by human activities.• Overland and streamflow travel time studies to be able to address possible spills response protocol and actions.• Enhance the continuous streamflow gauge network and update data regarding discharge to streams.• Update and verify outdated or missing water use data including Permit to Take Water (PTTW) information.• Development of acceptable water use targets to protect both the resource and the aquatic ecosystem.• Need for additional water quality monitoring sites.• Need for additional climatic sites/data monitoring.• Development of the ESRI ArcHydro data model. |

Table B-11: Data Identified

B1.7.2 Addressing Data

Future work will aim to develop and refine the overall database management system using the following methods:

- Preparing and refining land classification maps;
- Monitoring and analyzing groundwater level and chemistry involving both PGMN wells and municipal partner monitoring wells;
- Reviewing low-flow streamflow surveys (quality and quantity) to characterize discharge zones and associated water quality and to delineate zones that may be impacted by human activities;
- Reviewing overland and streamflow travel time studies to be able to address possible spills response protocol and actions;
- Enhancing the continuous streamflow gauge network and updating data regarding discharge to streams;
- Enhancing the coverage of climate data;
- Updating and verifying outdated or missing water using data including Permit to Take Water (PTTW) information; and
- Preparing a contaminant source database and associated risk to drinking water provided by each potential source.

Priority gaps that need to be addressed based on the analysis conducted include:

- Further development and promotion of its existing Clean Water Stewardship Program, which supports well upgrades and abandonment, nutrient management best management practices, and land restoration initiatives on private lands—all efforts that help remove potential pathways for contaminants;
- Need for additional water quality monitoring sites;
- Need for additional streamflow monitoring and climatic sites;
- Development of the ESRI ArcHydro data model; and
- Further estimates of water surplus (Thornwaite methodology).

B1.7.3 Assumptions

A number of statistical tests were performed on the surface water quality data for the study area. Tests were completed for all (PWQMN and CLOSPA) water quality sites that have a minimum of 40 samples (approximately five years of monitoring). Statistical tests were performed on chloride, nitrate, phosphorous, and copper.

Parametric and non-parametric statistical methods were used to describe the water data set and to determine if there are any temporal trends (i.e., the significance of a trend in water quality as it increases or decreases over time). Parametric statistical methods assume that observations are drawn from a normally distributed population. Since the distribution of surface water quality data are frequently skewed by extreme values, the assumptions of parametric statistical tests are often violated. Typically, non-parametric statistical methods are a more suitable choice for the analysis of surface water quality data. Non-parametric statistics do not assume a particular form of distribution, and they can handle outliers and non-detects that are common in water quality data.

The linear regression simply calculates a regression line on the time/value plot. A positive slope of the regression line indicates a trend towards increasing values, a negative slope indicates a trend to decreasing values. The results of this test should only be used qualitatively and should be confirmed by more sophisticated tests, such as Sen's and Mann-Kendall.

The Mann-Kendall test is a trend estimator that can be used to examine whether contaminant concentrations are diminishing or rising significantly over time. Unlike linear regression, there are no distributional assumptions. It is not greatly affected by gross data errors, outliers, or missing data (non-detects), and irregularly spaced measurement periods are allowed. Non-detects are assigned a value smaller than the smallest measured value.

The version of the Mann-Kendall Test used for this analysis comes from AquaChem and can be applied for virtually any surface water or groundwater parameter. The Mann Kendall test provides two values: S value and Z value. If the Mann-Kendall statistic (S) equals 0, then there is no increasing or decreasing trend in the data. If, however, S is less than 0, there is a decreasing trend, and if S is greater than 0, there is an increasing trend.

A two-sided test (for either increasing or decreasing trend) can also be obtained by using probability values. For monitoring stations with more than 10 samples, the normal approximation (Z) can be calculated. The quantity Z can be compared to standard normal cumulative distribution probabilities to test the null hypothesis of no trend.

B1.7.4 Method Limitations

Knowledge gaps relate to analysis and tool development to estimate and/or refine the water budget estimates and understand how the flow system operates. These tools enable us to predict the impact of potential future changes, such as increased municipal supply from groundwater due to climate change.

Priority knowledge gaps that need to be addressed include:

- Refinement of aquifer characterization and flow system understanding, including the orientation of bedrock valley systems and significant area recharge and discharge mapping;
- Development of surface water modelling capabilities;
- Refinement of a three-dimensional groundwater flow modelling tool;
- Refinement of the interaction of the surface water and groundwater flow models;
- Development of acceptable water use targets to protect both the resource and the aquatic ecosystem; and
- Development of methodology and tools to provide spills response analysis that will involve all pathways, including overland flow, stream travel, and groundwater flow, including the unsaturated zone transport.